

# Chapter 11

## Numerical Integration Methods in Vibration Analysis

$$11.1 \quad \frac{d^2 x}{dt^2} \Big|_i = \frac{\frac{dx}{dt} \Big|_{i+1} - \frac{dx}{dt} \Big|_i}{\Delta t} = \frac{\left(\frac{x_{i+2} - x_{i+1}}{\Delta t}\right) - \left(\frac{x_{i+1} - x_i}{\Delta t}\right)}{\Delta t} = \frac{x_{i+2} - 2x_{i+1} + x_i}{(\Delta t)^2}$$

$$\frac{d^3 x}{dt^3} \Big|_i = \frac{\frac{d^2 x}{dt^2} \Big|_{i+1} - \frac{d^2 x}{dt^2} \Big|_i}{\Delta t} = \frac{\left(\frac{x_{i+3} - 2x_{i+2} + x_{i+1}}{(\Delta t)^2}\right) - \left(\frac{x_{i+2} - 2x_{i+1} + x_i}{(\Delta t)^2}\right)}{\Delta t}$$

$$= \frac{x_{i+3} - 3x_{i+2} + 3x_{i+1} - x_i}{(\Delta t)^3}$$

$$\frac{d^4 x}{dt^4} \Big|_i = \frac{\frac{d^3 x}{dt^3} \Big|_{i+1} - \frac{d^3 x}{dt^3} \Big|_i}{\Delta t} = \frac{\left(\frac{x_{i+4} - 3x_{i+3} + 3x_{i+2} - x_{i+1}}{(\Delta t)^3}\right) - \left(\frac{x_{i+3} - 3x_{i+2} + 3x_{i+1} - x_i}{(\Delta t)^3}\right)}{\Delta t}$$

$$= \frac{x_{i+4} - 4x_{i+3} + 6x_{i+2} - 4x_{i+1} + x_i}{(\Delta t)^4}$$

$$11.2 \quad \frac{d^2 x}{dt^2} \Big|_i = \frac{\frac{dx}{dt} \Big|_i - \frac{dx}{dt} \Big|_{i-1}}{\Delta t} = \frac{\left(\frac{x_i - x_{i-1}}{\Delta t}\right) - \left(\frac{x_{i-1} - x_{i-2}}{\Delta t}\right)}{\Delta t} = \frac{x_i - 2x_{i-1} + x_{i-2}}{(\Delta t)^2}$$

$$\frac{d^3 x}{dt^3} \Big|_i = \frac{\frac{d^2 x}{dt^2} \Big|_i - \frac{d^2 x}{dt^2} \Big|_{i-1}}{\Delta t} = \frac{\left(\frac{x_i - 2x_{i-1} + x_{i-2}}{(\Delta t)^2}\right) - \left(\frac{x_{i-1} - 2x_{i-2} + x_{i-3}}{(\Delta t)^2}\right)}{\Delta t}$$

$$= \frac{x_i - 3x_{i-1} + 3x_{i-2} - x_{i-3}}{(\Delta t)^3}$$

$$\frac{d^4 x}{dt^4} \Big|_i = \frac{\frac{d^3 x}{dt^3} \Big|_i - \frac{d^3 x}{dt^3} \Big|_{i-1}}{\Delta t} = \frac{\left(\frac{x_i - 3x_{i-1} + 3x_{i-2} - x_{i-3}}{(\Delta t)^3}\right) - \left(\frac{x_{i-1} - 3x_{i-2} + 3x_{i-3} - x_{i-4}}{(\Delta t)^3}\right)}{\Delta t}$$

$$= \frac{x_i - 4x_{i-1} + 6x_{i-2} - 4x_{i-3} + x_{i-4}}{(\Delta t)^4}$$

$$11.3 \quad \frac{d^2 x}{dt^2} \Big|_i = \frac{x_{i+1} - 2x_i + x_{i-1}}{(\Delta t)^2}$$

$$\frac{d^4 x}{dt^4} \Big|_i = \frac{\frac{d^2 x}{dt^2} \Big|_{i+1} - 2 \frac{d^2 x}{dt^2} \Big|_i + \frac{d^2 x}{dt^2} \Big|_{i-1}}{(\Delta t)^2}$$

$$= \left\{ \left( \frac{x_{i+2} - 2x_{i+1} + x_i}{(\Delta t)^2} \right) - 2 \left( \frac{x_{i+1} - 2x_i + x_{i-1}}{(\Delta t)^2} \right) + \left( \frac{x_i - 2x_{i-1} + x_{i-2}}{(\Delta t)^2} \right) \right\} / (\Delta t)^2$$

$$= \frac{x_{i+2} - 4x_{i+1} + 6x_i - 4x_{i-1} + x_{i-2}}{(\Delta t)^4}$$

11.4 Equation:  $\ddot{x} + x = 0$   
 Central difference solution:  $x_{i+1} = (\Delta t)^2 \left\{ \left( \frac{2}{(\Delta t)^2} - 1 \right) x_i - \frac{1}{(\Delta t)^2} x_{i-1} \right\}$   
 $= (2 - \Delta t^2) x_i - x_{i-1}$  --- (E<sub>1</sub>)

For  $x_0 = 0$  and  $\dot{x}_0 = 1$ , Eq. (11.9) gives  $x_{-1} = -\Delta t$

(i) with  $\Delta t = 1$ ,  $x_{i+1} = x_i - x_{i-1}$  --- (E<sub>2</sub>)

Repetitive application of (E<sub>2</sub>), with  $x_{-1} = -1$  and  $x_0 = 0$ , gives the results shown in the following table.

(ii) With  $\Delta t = 0.5$ ,  $x_{i+1} = 1.75 x_i - x_{i-1}$  --- (E<sub>3</sub>)

The results, for  $x_{-1} = -0.5$  and  $x_0 = 0$ , are shown in the table.

comparison of solutions:

Time (t)	value of $x(t)$ obtained with		Exact value of $x(t)$ $x(t) = \sin t$
	$\Delta t = 1$	$\Delta t = 0.5$	
0	0	0	0
0.5	-	0.5	0.4794
1	1	0.8750	0.8415
1.5	-	1.0313	0.9975
2	1	0.9297	0.9093
2.5	-	0.5957	0.5985
3	0	0.1128	0.1411
3.5	-	-0.3983	-0.3508
4	-1	-0.8098	-0.7568
4.5	-	-1.0189	-0.9775
5	-1	-0.9733	-0.9589
5.5	-	-0.6843	-0.7055
6	0	-0.2242	-0.2794

11.5 At  $t_i$ , central difference formula gives

$$- \left( \frac{x_i - 2x_{i-1} + x_{i-2}}{(\Delta t)^2} \right) + 0.1 x_i = 0$$

i.e.  $x_i = \frac{2x_{i-1} - x_{i-2}}{1 - 0.1(\Delta t)^2} = 1.1111 (2x_{i-1} - x_{i-2})$  for  $\Delta t = 1$  --- (E<sub>1</sub>)

Since  $\dot{x}_0 = \frac{x_0 - x_{-1}}{\Delta t} = 0$ ,  $x_{-1} = x_0 = 1$

Eq. (E<sub>1</sub>) gives the following results.

$i$	1	2	3	4	5	6	7	8	9	10
$x_{i-1}$	1	1.1111	1.3580	1.7832	2.2084	2.6336	3.0588	3.4840	3.9092	4.3344
$x_{i-2}$	1	1	1.1111	1.3580	1.7832	2.2084	2.6336	3.0588	3.4840	3.9092
$x_i$ from (E <sub>1</sub> )	1.1111	1.3580	1.7832	2.2084	2.6336	3.0588	3.4840	3.9092	4.3344	4.7596

11.6

For given data, Eq. (11.7) gives  $x_{i+1} = 0.2(7x_i - 3x_{i-1})$  --- (E<sub>1</sub>)

Eqs. (11.8) and (11.9) give  $\ddot{x}_0 = -1$ ,  $x_{-1} = -0.625$

$$\left(\frac{c}{2m}\right)^2 = \left(\frac{1}{2}\right)^2 < \left(\frac{k}{m} = 1\right) \Rightarrow \text{Underdamped case.}$$

Since  $\tau_n = \frac{2\pi}{\omega_n} = 2\pi$  sec, we will consider the response for 15 steps.

$i+1$	$x_i$	$x_{i-1}$	$x_{i+1}$ from (E <sub>1</sub> )
1	0	-0.625	0.375
2	0.375	0	0.525
3	0.525000E+00	0.375000E+00	0.510000E+00
4	0.510000E+00	0.525000E+00	0.399000E+00
5	0.399000E+00	0.510000E+00	0.252600E+00
6	0.252600E+00	0.399000E+00	0.114240E+00
7	0.114240E+00	0.252600E+00	0.837604E-02
8	0.837604E-02	0.114240E+00	-0.568176E-01
9	-0.568176E-01	0.837604E-02	-0.845702E-01
10	-0.845702E-01	-0.568176E-01	-0.843078E-01
11	-0.843078E-01	-0.845702E-01	-0.672887E-01
12	-0.672887E-01	-0.843078E-01	-0.436196E-01
13	-0.436196E-01	-0.672887E-01	-0.206942E-01
14	-0.206942E-01	-0.436196E-01	-0.280008E-02
15	-0.280008E-02	-0.206942E-01	0.849639E-02

11.7

$\left(\frac{c}{2m}\right)^2 = 1 = \left(\frac{k}{m} = 1\right) \Rightarrow$  critically damped case.

For the data, Eqs. (11.7) to (11.9) give

$$x_{i+1} = \frac{1}{6}(7x_i - 2x_{i-1}) \quad \text{--- (E}_1\text{)}$$

$\ddot{x}_0 = -2$ ,  $x_{-1} = -0.75$ . Response is given below.

$i+1$	$x_i$	$x_{i-1}$	$x_{i+1}$ from (E <sub>1</sub> )
1	0	-0.75	0.25
2	0.25	0	0.2917
3	0.291667E+00	0.250000E+00	0.256944E+00
4	0.256944E+00	0.291667E+00	0.202546E+00
5	0.202546E+00	0.256944E+00	0.150656E+00
6	0.150656E+00	0.202546E+00	0.108250E+00
7	0.108250E+00	0.150656E+00	0.760728E-01
8	0.760728E-01	0.108250E+00	0.526683E-01
9	0.526683E-01	0.760728E-01	0.360888E-01

10	0.360888E-01	0.526683E-01	0.245475E-01
11	0.245475E-01	0.360888E-01	0.166091E-01
12	0.166091E-01	0.245475E-01	0.111948E-01
13	0.111948E-01	0.166091E-01	0.752425E-02
14	0.752425E-02	0.111948E-01	0.504668E-02
15	0.504668E-02	0.752425E-02	0.337971E-02

11.8 For given data, Eq. (11.7) gives  $x_{i+1} = 0.875 x_i$   
 This equation does not involve  $x_{i-1}$  and hence the response can't be found since  $x_0 = 0$ . Hence we change  $\Delta t$  to 0.4. This gives, from Eq. (11.7),  $x_{i+1} = 0.08889 (11.5 x_i - 1.25 x_{i-1})$  --- (E<sub>1</sub>)  
 Eqs. (11.8) and (11.9) give  $\ddot{x}_0 = -4$ ,  $x_{-1} = -0.72$

Results:

$i+1$	$x_{i-1}$	$x_i$	$x_{i+1}$ from (E <sub>1</sub> )
1	-0.720000E+00	0.000000E+00	0.800010E-01
2	0.000000E+00	0.800010E-01	0.817798E-01
3	0.800010E-01	0.817798E-01	0.747091E-01
4	0.817798E-01	0.747091E-01	0.672835E-01
5	0.747091E-01	0.672835E-01	0.604784E-01
6	0.672835E-01	0.604784E-01	0.543471E-01
7	0.604784E-01	0.543471E-01	0.488356E-01
8	0.543471E-01	0.488356E-01	0.438829E-01
9	0.488356E-01	0.438829E-01	0.394323E-01
10	0.438829E-01	0.394323E-01	0.354332E-01
11	0.394323E-01	0.354332E-01	0.318396E-01
12	0.354332E-01	0.318396E-01	0.286105E-01
13	0.318396E-01	0.286105E-01	0.257089E-01

11.9  $\omega_n = \sqrt{\frac{3000}{4}} = 27.3861$ ,  $\tau_n = 2\pi/\omega_n = 0.22943$ ;  $\Delta t = 0.05$   
 Eqs. (11.7) to (11.9) give

$$x_{i+1} = \frac{1}{1620} (200 x_i - 1580 x_{i-1} + F_i) \quad \text{where } F(t) = \begin{cases} 200 & ; 0 \leq t \leq 0.2 \\ -500t + 300 & ; 0.2 \leq t \leq 0.6 \end{cases}$$

--- (E<sub>1</sub>)

$\ddot{x}_0 = 50$ ,  $x_{-1} = 0.0625$

Results:

$i+1$	$t_i$	$F_i = F(t_i)$	$x_i$	$x_{i-1}$	$x_{i+1}$ from (E <sub>1</sub> )
1	0	0.200000E+03	0.000000E+00	0.625000E-01	0.625000E-01
2	0.05	0.200000E+03	0.625000E-01	0.000000E+00	0.131173E+00
3	0.1	0.200000E+03	0.131173E+00	0.625000E-01	0.786942E-01
4	0.15	0.200000E+03	0.786942E-01	0.131173E+00	0.523812E-02
5	0.2	0.200000E+03	0.523812E-02	0.786942E-01	0.473524E-01
6	0.25	0.175000E+03	0.473524E-01	0.523812E-02	0.108762E+00
7	0.3	0.150000E+03	0.108762E+00	0.473524E-01	0.598368E-01
8	0.35	0.125000E+03	0.598368E-01	0.108762E+00	-0.215287E-01
9	0.4	0.100000E+03	-0.215287E-01	0.598368E-01	0.711154E-03
10	0.45	0.750000E+02	0.711154E-03	-0.215287E-01	0.673812E-01
11	0.5	0.500000E+02	0.673812E-01	0.711154E-03	0.384892E-01
12	0.55	0.250000E+02	0.384892E-01	0.673812E-01	-0.455336E-01
13	0.6	-0.305176E-04	-0.455336E-01	0.384892E-01	-0.431603E-01

14	0.65	-0.250001E+02	-0.431603E-01	-0.455336E-01	0.236487E-01
15	0.7	-0.500001E+02	0.236487E-01	-0.431603E-01	0.141500E-01
16	0.75	-0.750001E+02	0.141500E-01	0.236487E-01	-0.676142E-01
17	0.8	-0.100000E+03	-0.676142E-01	0.141500E-01	-0.838765E-01
18	0.85	-0.125000E+03	-0.838765E-01	-0.676142E-01	-0.215709E-01
19	0.9	-0.150000E+03	-0.215709E-01	-0.838765E-01	-0.134502E-01
20	0.95	-0.175000E+03	-0.134502E-01	-0.215709E-01	-0.886470E-01
21	1.00	-0.200000E+03	-0.886470E-01	-0.134502E-01	-0.121283E+00

11.10 Equation:  $m\ddot{x} + c\dot{x} + kx = F(t) = t$

Since  $\tau_n = 2\pi$ ,  $\Delta t$  is selected as 0.5.

Eqs. (11.7) to (11.9) give  $x_{i+1} = 0.2(7x_i - 3x_{i-1} + t_i) \dots (E_1)$

$\ddot{x}_0 = 0, x_{-1} = 0$

Exact solution (from Example 4.9):

For  $\delta F = 1, k = 1, \omega_n = 1, \gamma = 0.5, \omega_d = 0.86603$ , we find

$x(t) = t - 1 + e^{-0.5t} (\cos 0.86603t - 0.57735 \sin 0.86603t) \dots (E_2)$

Results:

$i+1$	$t_i$	$x_i$	$x_{i-1}$	$x_{i+1}, (E_1)$	$x_{i+1}, (E_2)$
1	0	0.000000E+00	0.000000E+00	0.000000E+00	0.182477E-01
2	0.5	0.000000E+00	0.000000E+00	0.100000E+00	0.126190E+00
3	1	0.100000E+00	0.000000E+00	0.340000E+00	0.364080E+00
4	1.5	0.340000E+00	0.100000E+00	0.716000E+00	0.731292E+00
5	2	0.716000E+00	0.340000E+00	0.119840E+01	0.120253E+01
6	2.5	0.119840E+01	0.716000E+00	0.174816E+01	0.174240E+01
7	3	0.174816E+01	0.119840E+01	0.232838E+01	0.231622E+01
8	3.5	0.232838E+01	0.174816E+01	0.291084E+01	0.289641E+01
9	4	0.291084E+01	0.232838E+01	0.347815E+01	0.346501E+01
10	4.5	0.347815E+01	0.291084E+01	0.402290E+01	0.401335E+01
11	5	0.402290E+01	0.347815E+01	0.454518E+01	0.454011E+01
12	5.5	0.454518E+01	0.402290E+01	0.504950E+01	0.504860E+01
13	6	0.504950E+01	0.454518E+01	0.554220E+01	0.554439E+01
14	6.5	0.554220E+01	0.504950E+01	0.602938E+01	0.603328E+01
15	7	0.602938E+01	0.554220E+01	0.651581E+01	0.652013E+01
16	7.5	0.651581E+01	0.602938E+01	0.700451E+01	0.700828E+01
17	8	0.700451E+01	0.651581E+01	0.749682E+01	0.749949E+01
18	8.5	0.749682E+01	0.700451E+01	0.799285E+01	0.799426E+01
19	9	0.799285E+01	0.749682E+01	0.849190E+01	0.849219E+01
20	9.5	0.849190E+01	0.799285E+01	0.899294E+01	0.899244E+01

11.11 Let  $x_1 = x, x_2 = \frac{dx_1}{dt} = \frac{dx}{dt}, x_3 = \frac{dx_2}{dt} = \frac{d^2x}{dt^2}, \dots, x_n = \frac{dx_{n-1}}{dt} = \frac{d^{n-1}x}{dt^{n-1}}$

Equation:  $\frac{d^n x}{dt^n} = \frac{g(x,t)}{a_n} - \frac{a_{n-1}}{a_n} \frac{d^{n-1}x}{dt^{n-1}} - \dots - \frac{a_1}{a_n} \frac{dx}{dt} \dots (E_1)$

$= \frac{g(x,t)}{a_n} - \frac{a_{n-1}}{a_n} x_n - \frac{a_{n-2}}{a_n} x_{n-1} - \dots - \frac{a_1}{a_n} x_2 \dots (E_2)$

(E<sub>1</sub>) and (E<sub>2</sub>) can be expressed as

$\frac{d\vec{X}}{dt} = \vec{F}(\vec{X}, t)$

where  $\vec{X} = \begin{Bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{Bmatrix}$  and  $\vec{F} = \begin{Bmatrix} f_1 \\ f_2 \\ f_3 \\ \vdots \\ f_n \end{Bmatrix} = \begin{Bmatrix} x_2 \\ x_3 \\ x_4 \\ \vdots \\ x_n \\ \frac{g(x,t)}{a_n} - \frac{a_{n-1}}{a_n} x_n - \dots - \frac{a_1}{a_n} x_2 \end{Bmatrix}$

11.12 The main program for problems (a) and (b) is given below.

```

=====
C MAIN PROGRAM FOR CALLING THE SUBROUTINE RK4
C =====
C THE FOLLOWING 7 LINES NEED TO BE CHANGED FOR A DIFFERENT PROBLEM
  DIMENSION TIME(40),X(40,1),XX(1),F(1),YI(1),YJ(1),YK(1),YL(1),
2  UU(1)
  XX(1)=1.0
  NEQ=1
  NSTEP=40
  DT=0.1
  T=0.0
  WRITE (58,10)
10  FORMAT (//,3X,5H I ,10H TIME(1),7X,5H X(1),/)
  DO 40 I=1,NSTEP
  CALL RK4 (T,DT,NEQ,XX,F,YI,YJ,YK,YL,UU)
  TIME(I)=T
  DO 20 J=1,NEQ
20  X(I,J)=XX(J)
  WRITE (58,30) I,TIME(I),(X(I,J),J=1,NEQ)
30  FORMAT (2X,15,F10.4,2X,E15.8,2X,E15.8)
40  CONTINUE
  STOP
  END

```

The subroutine FUN and output are given below.

For problem (a)			For problem (b)		
SUBROUTINE FUN (X,F,N,I)			SUBROUTINE FUN (X,F,N,I)		
DIMENSION X(N),F(N)			DIMENSION X(N),F(N)		
F(1)=X(1)-1.5*EXP(-0.5*T)			F(1)=-T*(X(1)**2)		
RETURN			RETURN		
END			END		
I	TIME(I)	X(I)	I	TIME(I)	X(I)
1	0.1000	0.95122939E+00	1	0.1000	0.99502486E+00
2	0.2000	0.90483737E+00	2	0.2000	0.98039210E+00
3	0.3000	0.86070788E+00	3	0.3000	0.95693767E+00
4	0.4000	0.81873059E+00	4	0.4000	0.92592573E+00
5	0.5000	0.77880061E+00	5	0.5000	0.88888866E+00
6	0.6000	0.74081802E+00	6	0.6000	0.84745741E+00
7	0.7000	0.70468783E+00	7	0.7000	0.80321264E+00
8	0.8000	0.67031974E+00	8	0.8000	0.75757557E+00

31	3.1000	0.21224274E+00	31	3.1000	0.17226547E+00
32	3.2000	0.20189069E+00	32	3.2000	0.16339886E+00
33	3.3000	0.19204344E+00	33	3.3000	0.15515921E+00
34	3.4000	0.18267635E+00	34	3.4000	0.14749278E+00
35	3.5000	0.17376597E+00	35	3.5000	0.14035103E+00
36	3.6000	0.16529004E+00	36	3.6000	0.13368998E+00
37	3.7000	0.15722735E+00	37	3.7000	0.12746987E+00
38	3.8000	0.14955774E+00	38	3.8000	0.12165464E+00
39	3.9000	0.14226201E+00	39	3.9000	0.11621163E+00
40	4.0000	0.13532193E+00	40	4.0000	0.11111123E+00

11.13

The program and output are given below.

```

C =====
C
C MAIN PROGRAM FOR CALLING THE SUBROUTINE RK2
C
C =====
C THE FOLLOWING 6 LINES NEED TO BE CHANGED FOR A DIFFERENT PROBLEM
  DIMENSION TIME(40),X(40,2),XX(2),F(2),YI(2),YJ(2),UU(2)
  XX(1)=0.0
  XX(2)=0.0
  NEQ=2
  NSTEP=40
  DT=0.31416/2
  I=0.0
  WRITE (57,10)
10  FORMAT (//,3X,5H I ,10H TIME(I),7X,5H X(1),12X,5H X(2),/)
  DO 40 I=1,NSTEP
  CALL RK2 (T,DT,NEQ,XX,F,YI,YJ,UU)
  TIME(I)=T
  DO 20 J=1,NEQ
20  X(I,J)=XX(J)
  WRITE (57,30) I,TIME(I),(X(I,J),J=1,NEQ)
30  FORMAT (2X,I5,F10.4,2X,E15.8,2X,E15.8)
40  CONTINUE
  STOP
  END
C =====
C SUBROUTINE RK2
C =====
  SUBROUTINE RK2 (T,DT,N,XX,F,XI,XJ,UU)
  DIMENSION XI(N),XJ(N),UU(N),XX(N),F(N)
  DO 10 I=1,N
10  UU(I)=XX(I)
  CALL FUN (XX,F,N,T)
  DO 20 I=1,N
  XI(I)=F(I)*DT
20  XX(I)=UU(I)+XI(I)
  T=T+DT
  CALL FUN (XX,F,N,T)
  DO 30 I=1,N
  XJ(I)=F(I)*DT
30  XX(I)=UU(I)+(XI(I)+XJ(I))/2.0
  RETURN
  END

```

```

C =====
C SUBROUTINE FUN FOR USE IN THE SUBROUTINE RK2
C THIS SUBROUTINE CHANGES FROM PROBLEM TO PROBLEM
C =====
      SUBROUTINE FUN (X,F,N,T)
      DIMENSION X(N),F(N)
      F(1)=X(2)
      F0=1.0
      T0=3.1416
      XM=1.0
      XC=0.2
      XK=1.0
      FT=F0*(1.0-SIN(3.1416*T/(2.0*T0)))
      F(2)=(FT-XC*X(2)-XK*X(1))
      RETURN
      END

```

I	TIME(I)	X(1)	X(2)
1	0.1571	0.12337063E-01	0.14845039E+00
2	0.3142	0.46506263E-01	0.27647862E+00
3	0.4712	0.99086702E-01	0.38170516E+00
4	0.6283	0.16633771E+00	0.46245623E+00
5	0.7854	0.24431184E+00	0.51778144E+00
6	0.9425	0.32896915E+00	0.54745305E+00
7	1.0996	0.41628993E+00	0.55194682E+00
8	1.2566	0.50238299E+00	0.53240609E+00
9	1.4137	0.58358723E+00	0.49059010E+00
10	1.5708	0.65656364E+00	0.42880845E+00
:			
36	5.6549	-0.28463572E+00	0.51887971E+00
37	5.8120	-0.19237404E+00	0.65321267E+00
38	5.9690	-0.79548687E-01	0.77926701E+00
39	6.1261	0.52324414E-01	0.89440674E+00
40	6.2832	0.20133464E+00	0.99627036E+00

11.14 The computer program and output are given below.

```

C =====
C
C MAIN PROGRAM FOR CALLING THE SUBROUTINE RK3
C
C =====
C THE FOLLOWING 6 LINES NEED TO BE CHANGED FOR A DIFFERENT PROBLEM
      DIMENSION TIME(40),X(40,2),XX(2),F(2),YI(2),YJ(2),YK(2),UU(2)
      XX(1)=0.0
      XX(2)=0.0
      NEQ=2
      NSTEP=40
      DT=0.31416/2
      T=0.0
      WRITE (57,10)
10  FORMAT (//,3X,5H I ,10H TIME(I),7X,5H X(1),12X,5H X(2),/)
      DO 40 I=1,NSTEP
      CALL RK3 (T,DT,NEQ,XX,F,YI,YJ,YK,UU)
      TIME(I)=T

```

```

      DO 20 J=1,NEQ
20    X(I,J)=XX(J)
      WRITE (57,30) I,TIME(I),(X(I,J),J=1,NEQ)
30    FORMAT (2X,I5,F10.4,2X,E15.8,2X,E15.8)
40    CONTINUE
      STOP
      END

```

```

C =====
C SUBROUTINE RK3

```

```

C =====
      SUBROUTINE RK3 (T,DT,N,XX,F,XI,XJ,XK,UU)
      DIMENSION XI(N),XJ(N),XK(N),UU(N),XX(N),F(N)
      DO 10 I=1,N
10    UU(I)=XX(I)
      CALL FUN (XX,F,N,T)
      DO 20 I=1,N
20    XI(I)=F(I)*DT
      XX(I)=UU(I)+XI(I)/2.0
      T=T+DT/2.0
      CALL FUN (XX,F,N,T)
      DO 30 I=1,N
30    XJ(I)=F(I)*DT
      XX(I)=UU(I)-XI(I)+2.0*XJ(I)
      CALL FUN (XX,F,N,T)
      DO 40 I=1,N
40    XK(I)=F(I)*DT
      XX(I)=UU(I)+(XI(I)+4.0*XJ(I)+XK(I))/6.0
      RETURN
      END

```

```

C =====
C SUBROUTINE FUN FOR USE IN THE SUBROUTINE RK3
C THIS SUBROUTINE CHANGES FROM PROBLEM TO PROBLEM
C =====

```

```

      SUBROUTINE FUN (X,F,N,T)
      DIMENSION X(N),F(N)
      F(1)=X(2)
      F0=1.0
      T0=3.1416
      XM=1.0
      XC=0.2
      XK=1.0
      FT=F0*(1.0-SIN(3.1416*T/(2.0*T0)))
      F(2)=(FT-XC*X(2)-XK*X(1))
      RETURN
      END

```

I	TIME(I)	X(1)	X(2)
1	0.0785	0.11884968E-01	0.14891791E+00
2	0.1571	0.46078302E-01	0.28356332E+00
3	0.2356	0.10011104E+00	0.40113181E+00
4	0.3142	0.17111324E+00	0.49932912E+00
5	0.3927	0.25589743E+00	0.57641083E+00
6	0.4712	0.35104716E+00	0.63120759E+00
7	0.5498	0.45300832E+00	0.66313553E+00
8	0.6283	0.55818105E+00	0.67219222E+00
9	0.7069	0.66300964E+00	0.65893871E+00
10	0.7854	0.76406908E+00	0.62446821E+00

36	2.8274	-0.56948423E+00	-0.28658101E+00
37	2.9060	-0.60655731E+00	-0.18535951E+00
38	2.9845	-0.62764353E+00	-0.83412744E-01
39	3.0631	-0.63280767E+00	0.17002784E-01
40	3.1416	-0.62246108E+00	0.11373823E+00

11.15 The main program, subroutine FUN and results are given.

```

C =====
C
C MAIN PROGRAM FOR CALLING THE SUBROUTINE RK2
C
C =====
C THE FOLLOWING 6 LINES NEED TO BE CHANGED FOR A DIFFERENT PROBLEM
  DIMENSION TIME(40),X(40,2),XX(2),F(2),YI(2),YJ(2),UU(2)
  XX(1)=5.0
  XX(2)=0.0
  NEQ=2
  NSTEP=40
  DT=0.01
  T=0.0
  WRITE (61,10)
10  FORMAT (//,3X,5H I ,10H  TIME(1),7X,5H X(1),12X,5H X(2),/)
  DO 40 I=1,NSTEP
  CALL RK2 (T,DT,NEQ,XX,F,YI,YJ,UU)
  TIME(I)=T
  DO 20 J=1,NEQ
20  X(1,J)=XX(J)
  WRITE (61,30) I,TIME(I),(X(1,J),J=1,NEQ)
30  FORMAT (2X,I5,F10.4,2X,E15.8,2X,E15.8)
40  CONTINUE
  STOP
  END

SUBROUTINE FUN (X,F,N,T)
  DIMENSION X(N),F(N)
  F(1)=X(2)
  F(2)=-1000.0*X(1)
  RETURN
  END

```

I	TIME(I)	X(1)	X(2)
1	0.0100	0.47500000E+01	-0.50000000E+02
2	0.0200	0.40124998E+01	-0.95000000E+02
3	0.0300	0.28618748E+01	-0.13037500E+03
4	0.0400	0.14150311E+01	-0.15247501E+03
5	0.0500	-0.18047059E+00	-0.15900157E+03
6	0.0600	-0.17614628E+01	-0.14924678E+03
7	0.0700	-0.31658576E+01	-0.12416982E+03
8	0.0800	-0.42492628E+01	-0.86302750E+02
9	0.0900	-0.48998270E+01	-0.39494984E+02
10	0.1000	-0.50497856E+01	0.11478039E+02
:			

36	0.3600	0.28330812E+01	0.13901421E+03
37	0.3700	0.40815692E+01	0.10373269E+03
38	0.3800	0.49148178E+01	0.57730366E+02
39	0.3900	0.52463808E+01	0.56956673E+01
40	0.4000	0.50410185E+01	-0.47052921E+02

11.16

The main program, subroutine FUN and results are given.

```

C =====
C
C MAIN PROGRAM FOR CALLING THE SUBROUTINE RK3
C
C =====
C THE FOLLOWING 6 LINES NEED TO BE CHANGED FOR A DIFFERENT PROBLEM
  DIMENSION TIME(40),X(40,2),XX(2),F(2),YI(2),YJ(2),YK(2),UU(2)
  XX(1)=5.0
  XX(2)=0.0
  NEQ=2
  NSTEP=40
  DT=0.01
  T=0.0
  WRITE (62,10)
10  FORMAT (//,3X,5H I ,10H  TIME(I),7X,5H X(1),12X,5H X(2),/)
  DO 10 I=1,NSTEP
  CALL RK3 (T,DT,NEQ,XX,F,YI,YJ,YK,UU)
  TIME(I)=T
  DO 20 J=1,NEQ
20  X(I,J)=XX(J)
  WRITE (62,30) I,TIME(I),(X(I,J),J=1,NEQ)
30  FORMAT (2X,I5,F10.4,2X,E15.8,2X,E15.8)
40  CONTINUE
  STOP
  END

  SUBROUTINE FUN (X,F,N,T)
  DIMENSION X(N),F(N)
  F(1)=X(2)
  F(2)=-1000.0*X(1)
  RETURN
  END

```

I	TIME(I)	X(1)	X(2)
1	0.0050	0.47500000E+01	-0.49166668E+02
2	0.0100	0.40290279E+01	-0.93416672E+02
3	0.0150	0.29089794E+01	-0.12836461E+03
4	0.0200	0.15012785E+01	-0.15055135E+03
5	0.0250	-0.54207087E-01	-0.15778635E+03
6	0.0300	-0.16030624E+01	-0.14936400E+03
7	0.0350	-0.29916553E+01	-0.12613235E+03
8	0.0400	-0.40823741E+01	-0.90407799E+02
9	0.0450	-0.47672653E+01	-0.45744061E+02
10	0.0500	-0.49787188E+01	0.34212494E+01
⋮			
36	0.1800	0.18843936E+01	0.14399376E+03
37	0.1850	0.32061126E+01	0.11826420E+03
38	0.1900	0.42087383E+01	0.80824219E+02
39	0.1950	0.47930727E+01	0.35397079E+02
40	0.2000	0.49014902E+01	-0.13504654E+02

11.17 The main program, subroutine FUN and output are given.

```

C =====
C
C MAIN PROGRAM FOR CALLING THE SUBROUTINE RK4
C
C =====
C THE FOLLOWING 7 LINES NEED TO BE CHANGED FOR A DIFFERENT PROBLEM
  DIMENSION TIME(40),X(40,2),XX(2),F(2),YI(2),YJ(2),YK(2),YL(2),
2  UU(2)
  XX(1)=5.0
  XX(2)=0.0
  NEQ=2
  NSTEP=40
  DT=0.01
  T=0.0
  WRITE (59,10)
10  FORMAT (//,3X,5H I ,10H TIME(I),7X,5H X(1),12X,5H X(2),/)
  DO 40 I=1,NSTEP
  CALL RK4 (T,DT,NEQ,XX,F,YI,YJ,YK,YL,UU)
  TIME(I)=T
  DO 20 J=1,NEQ
20  X(I,J)=XX(J)
  WRITE (59,30) I,TIME(I),(X(I,J),J=1,NEQ)
30  FORMAT (2X,I5,F10.4,2X,E15.8,2X,E15.8)
40  CONTINUE
  STOP
  END

SUBROUTINE FUN (X,F,N,T)
DIMENSION X(N),F(N)
F(1)=X(2)
F(2)=-1000.0*X(1)
RETURN
END

```

I	TIME(I)	X(1)	X(2)
1	0.0100	0.47520833E+01	-0.49166668E+02
2	0.0200	0.40329871E+01	-0.93457642E+02
3	0.0300	0.29140182E+01	-0.12848141E+03
4	0.0400	0.15061308E+01	-0.15076540E+03
5	0.0500	-0.51074505E-01	-0.15810023E+03
6	0.0600	-0.16031944E+01	-0.14975887E+03
7	0.0700	-0.29963315E+01	-0.12656857E+03
8	0.0800	-0.40923543E+01	-0.90828957E+02
9	0.0900	-0.47825933E+01	-0.46083870E+02
10	0.1000	-0.49986143E+01	0.32299538E+01
:			
36	0.3600	0.18898817E+01	0.14634213E+03
37	0.3700	0.32352061E+01	0.12050217E+03
38	0.3800	0.42597318E+01	0.82714409E+02
39	0.3900	0.48618784E+01	0.36725792E+02
40	0.4000	0.49819474E+01	-0.12903667E+02

11.18 The main program, subroutine EXTFUN and results are given.

```

C =====
C
C PROGRAM -
C MAIN PROGRAM WHICH CALLS CDIFF
C
C =====
C FOLLOWING 10 LINES CONTAIN PROBLEM-DEPENDENT DATA
  REAL M(2,2),K(2,2),MC(2,2),MK(2,2),MCI(2,2),MMC(2,2)
  DIMENSION C(2,2),XI(2),XDI(2),XDDI(2),XM1(2),F(2),R(2),RR(2),
2  XMK(2),XMI(2),XM2(2),XP1(2),ZA(2),ZB(2),ZC(2),LA(2),LB(2,2),
3  S(2),X(25,2),XD(25,2),XDD(25,2)
  DATA N,NSTEP,NSTEP1,DELT/2,24,25,0.24216267/
  DATA XI/0.0,0.0/
  DATA XDI/0.0,0.0/
  DATA M/1.0,0.0,0.0,2.0/
  DATA C/2.0,-2.0,-2.0,2.0/
  DATA K/6.0,-2.0,-2.0,8.0/
C END OF PROBLEM-DEPENDENT DATA
  CALL CDIFF (M,C,K,XI,XDI,XDDI,N,NSTEP,DELT,F,R,RR,XM1,XM2,XP1,
2  MC,MK,MCI,XMK,MMC,XMI,ZA,ZB,ZC,LA,LB,S,X,XD,XDD,NSTEP1)
  WRITE (53,10)
10  FORMAT (//,38H SOLUTION BY CENTRAL DIFFERENCE METHOD,/)
  WRITE (53,20) N,NSTEP,DELT
20  FORMAT (12H GIVEN DATA:,//,3H N=,I5,4X,7H NSTEP=,I5,4X,6H DELT=,
2  E15.8,/)
  WRITE (53,30)
30  FORMAT (10H SOLUTION:,//,5H STEP,3X,5H TIME,3X,7H X(I,1),3X,
2  8H XD(I,1),2X,9H XDD(I,1),4X,7H X(I,2),3X,8H XD(I,2),2X,
3  9H XDD(I,2),/)
  DO 40 I=1,NSTEP1
  TIME=REAL(I-1)*DELT
40  WRITE (53,50) I,TIME,X(I,1),XD(I,1),XDD(I,1),X(I,2),XD(I,2),
2  XDD(I,2)
50  FORMAT (1X,I4,F8.4,6(1X,E10.4))
  STOP
  END
C =====
C
C SUBROUTINE EXTFUN
C THIS SUBROUTINE IS PROBLEM-DEPENDENT
C
C =====
  SUBROUTINE EXTFUN (F,TIME,N)
  DIMENSION F(N)
  F(1)=0.0
  F(2)=10.0
  RETURN
  END
  SOLUTION BY CENTRAL DIFFERENCE METHOD
  GIVEN DATA:
  N=      2      NSTEP=    24      DELT= 0.24216267E+00

```

## SOLUTION:

STEP	TIME	X(I,1)	XD(I,1)	XDD(I,1)	X(I,2)	XD(I,2)	XDD(I,2)
1	0.0000	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.5000E+01
2	0.2422	0.0000E+00	0.0000E+00	0.0000E+00	0.1466E+00	0.3077E-07	0.5000E+01
3	0.4843	0.1122E+00	0.2317E+00	0.1913E+01	0.5045E+00	0.1042E+01	0.3604E+01
4	0.7265	0.3601E+00	0.7436E+00	0.2314E+01	0.9859E+00	0.1733E+01	0.2105E+01
5	0.9687	0.6966E+00	0.1207E+01	0.1510E+01	0.1500E+01	0.2056E+01	0.5669E+00
6	1.2108	0.1036E+01	0.1396E+01	0.5531E-01	0.1961E+01	0.2013E+01	-.9224E+00
7	1.4530	0.1289E+01	0.1224E+01	-.1478E+01	0.2292E+01	0.1633E+01	-.2217E+01
8	1.6951	0.1388E+01	0.7260E+00	-.2635E+01	0.2438E+01	0.9851E+00	-.3136E+01
9	1.9373	0.1302E+01	0.2610E-01	-.3145E+01	0.2378E+01	0.1795E+00	-.3518E+01
10	2.1795	0.1045E+01	-.7088E+00	-.2925E+01	0.2127E+01	-.6433E+00	-.3277E+01
11	2.4216	0.6664E+00	-.1312E+01	-.2061E+01	0.1732E+01	-.1335E+01	-.2439E+01
12	2.6638	0.2433E+00	-.1655E+01	-.7648E+00	0.1270E+01	-.1769E+01	-.1146E+01
13	2.9060	-.1398E+00	-.1665E+01	0.6808E+00	0.8289E+00	-.1864E+01	0.3640E+00
14	3.1481	-.4070E+00	-.1343E+01	0.1979E+01	0.4940E+00	-.1602E+01	0.1803E+01
15	3.3903	-.5055E+00	-.7550E+00	0.2874E+01	0.3288E+00	-.1033E+01	0.2895E+01
16	3.6324	-.4166E+00	-.2001E-01	0.3196E+01	0.3645E+00	-.2673E+00	0.3427E+01
17	3.8746	-.1584E+00	0.7167E+00	0.2889E+01	0.5935E+00	0.5466E+00	0.3295E+01
18	4.1168	0.2183E+00	0.1311E+01	0.2019E+01	0.9707E+00	0.1252E+01	0.2527E+01
19	4.3589	0.6395E+00	0.1647E+01	0.7600E+00	0.1422E+01	0.1712E+01	0.1271E+01
20	4.6011	0.1023E+01	0.1662E+01	-.6399E+00	0.1861E+01	0.1838E+01	-.2264E+00
21	4.8433	0.1295E+01	0.1353E+01	-.1908E+01	0.2201E+01	0.1608E+01	-.1675E+01
22	5.0854	0.1403E+01	0.7832E+00	-.2800E+01	0.2378E+01	0.1067E+01	-.2793E+01
23	5.3276	0.1326E+01	0.6342E-01	-.3145E+01	0.2357E+01	0.3211E+00	-.3366E+01
24	5.5697	0.1080E+01	-.6657E+00	-.2877E+01	0.2143E+01	-.4839E+00	-.3283E+01
25	5.8119	0.7142E+00	-.1263E+01	-.2053E+01	0.1779E+01	-.1192E+01	-.2564E+01

11.19 The problem-dependent data to be used in the main program (which calls CDIFF), subroutine EXTFUN and output are given below.

C FOLLOWING 10 LINES CONTAIN PROBLEM-DEPENDENT DATA

```

REAL M(2,2),K(2,2),MC(2,2),MK(2,2),MCI(2,2),MMC(2,2)
DIMENSION C(2,2),XI(2),XDI(2),XDDI(2),XM1(2),F(2),R(2),RR(2),
2 XMK(2),XMI(2),XM2(2),XP1(2),ZA(2),ZB(2),ZC(2),LA(2),LB(2,2),
3 S(2),X(25,2),XD(25,2),XDD(25,2)
DATA N,NSTEP,NSTEP1,DELT/2,24,25,0.24216267/
DATA XI/0.0,0.0/
DATA XDI/0.0,0.0/
DATA M/1.0,0.0,0.0,2.0/
DATA C/0.0,0.0,0.0,0.0/
DATA K/6.0,-2.0,-2.0,8.0/

```

C END OF PROBLEM-DEPENDENT DATA

```

SUBROUTINE EXTFUN (F,TIME,N)
DIMENSION F(N)
F(1)=10.0*SIN(5.0*TIME)
F(2)=0.0
RETURN
END

```

SOLUTION BY CENTRAL DIFFERENCE METHOD

GIVEN DATA:

N= 2 NSTEP= 24 DELT= 0.24216267E+00

SOLUTION:

STEP	TIME	X(I,1)	XD(I,1)	XDD(I,1)	X(I,2)	XD(I,2)	XDD(I,2)
1	0.0000	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00
2	0.2422	0.5488E+00	0.1133E+01	0.9359E+01	0.0000E+00	0.0000E+00	0.0000E+00
3	0.4843	0.1291E+01	0.2666E+01	0.3301E+01	0.3219E-01	0.6645E-01	0.5488E+00
4	0.7265	0.1307E+01	0.1565E+01	-.1240E+02	0.1325E+00	0.2737E+00	0.1162E+01
5	0.9687	0.2964E+00	-.2054E+01	-.1749E+02	0.2784E+00	0.5084E+00	0.7765E+00
6	1.2108	-.9186E+00	-.4595E+01	-.3493E+01	0.3764E+00	0.5035E+00	-.8173E+00
7	1.4530	-.1279E+01	-.3252E+01	0.1458E+02	0.3322E+00	0.1110E+00	-.2424E+01
8	1.6951	-.6732E+00	0.5066E+00	0.1647E+02	0.1351E+00	-.4982E+00	-.2608E+01
9	1.9373	0.3324E-01	0.2709E+01	0.1722E+01	-.1332E+00	-.9609E+00	-.1214E+01
10	2.1795	0.1288E+00	0.1656E+01	-.1042E+02	-.3683E+00	-.1039E+01	0.5660E+00
11	2.4216	-.1236E+00	-.3238E+00	-.5933E+01	-.5094E+00	-.7768E+00	0.1602E+01
12	2.6638	0.8610E-02	-.2481E+00	0.6558E+01	-.5383E+00	-.3511E+00	0.1914E+01
13	2.9060	0.6165E+00	0.1528E+01	0.8111E+01	-.4404E+00	0.1424E+00	0.2162E+01
14	3.1481	0.9366E+00	0.1916E+01	-.4906E+01	-.2031E+00	0.6921E+00	0.2378E+01
15	3.3903	0.3481E+00	-.5540E+00	-.1549E+02	0.1368E+00	0.1192E+01	0.1749E+01
16	3.6324	-.7189E+00	-.3418E+01	-.8160E+01	0.4650E+00	0.1380E+01	-.1991E+00
17	3.8746	-.1185E+01	-.3166E+01	0.1024E+02	0.6420E+00	0.1043E+01	-.2579E+01
18	4.1168	-.5807E+00	0.2853E+00	0.1826E+02	0.5989E+00	0.2764E+00	-.3753E+01
19	4.3589	0.4129E+00	0.3300E+01	0.6635E+01	0.3812E+00	-.5385E+00	-.2976E+01
20	4.6011	0.8081E+00	0.2867E+01	-.1021E+02	0.9837E-01	-.1033E+01	-.1112E+01
21	4.8433	0.4651E+00	0.1077E+00	-.1259E+02	-.1602E+00	-.1118E+01	0.4146E+00
22	5.0854	0.1098E+00	-.1442E+01	-.2098E+00	-.3539E+00	-.9338E+00	0.1106E+01
23	5.3276	0.2596E+00	-.4244E+00	0.8612E+01	-.4582E+00	-.6152E+00	0.1525E+01
24	5.5697	0.5064E+00	0.8188E+00	0.1655E+01	-.4397E+00	-.1772E+00	0.2092E+01
25	5.8119	0.1089E+00	-.3112E+00	-.1099E+02	-.2885E+00	0.3504E+00	0.2265E+01

11.20 The problem-dependent data to be used in the main program (which calls CDIFF), subroutine EXTFUN and output are given below.

```

C FOLLOWING 10 LINES CONTAIN PROBLEM-DEPENDENT DATA
  REAL M(2,2),K(2,2),MC(2,2),MK(2,2),MCI(2,2),MMC(2,2)
  DIMENSION C(2,2),XI(2),XDI(2),XDD1(2),XM1(2),F(2),R(2),RR(2),
  2  XMK(2),XM2(2),XP1(2),ZA(2),ZB(2),ZC(2),LA(2),LB(2,2),
  3  S(2),X(25,2),XD(25,2),XDD(25,2)
  DATA N,NSTEP,NSTEP1,DELT/2,24,25,0.25/
  DATA XI/0.0,0.0/
  DATA XDI/0.0,0.0/
  DATA M/2.0,0.0,0.0,1.0/
  DATA C/0.0,0.0,0.0,0.0/
  DATA K/6.0,-2.0,-2.0,4.0/
C END OF PROBLEM-DEPENDENT DATA
SUBROUTINE EXTFUN (F,TIME,N)
  DIMENSION F(N)
  F(1)=5.0
  F(2)=20.0*SIN(5.0*TIME)
  RETURN
END

```

SOLUTION BY CENTRAL DIFFERENCE METHOD

GIVEN DATA:


N= 2      NSTEP= 24      DELT= 0.25000000E+00

SOLUTION:

STEP	TIME	X(I,1)	XD(I,1)	XDD(I,1)	X(I,2)	XD(I,2)	XDD(I,2)
------	------	--------	---------	----------	--------	---------	----------

1	0.0000	0.0000E+00	0.0000E+00	0.2500E+01	0.0000E+00	0.0000E+00	0.0000E+00
2	0.2500	0.7813E-01	0.0000E+00	0.2500E+01	0.1186E+01	0.2372E+01	0.1898E+02
3	0.5000	0.3720E+00	0.7440E+00	0.3452E+01	0.2834E+01	0.5668E+01	0.7381E+01
4	0.7500	0.9295E+00	0.1703E+01	0.4218E+01	0.3105E+01	0.3837E+01	-.2202E+02
5	1.0000	0.1663E+01	0.2582E+01	0.2816E+01	0.1517E+01	-.2633E+01	-.2974E+02
6	1.2500	0.2336E+01	0.2813E+01	-.9716E+00	-.2832E+00	-.6776E+01	-.3407E+01
7	1.5000	0.2709E+01	0.2092E+01	-.4790E+01	-.5484E+00	-.4131E+01	0.2456E+02
8	1.7500	0.2697E+01	0.7215E+00	-.6176E+01	0.4430E+00	0.1452E+01	0.2011E+02
9	2.0000	0.2362E+01	-.6938E+00	-.5147E+01	0.9807E+00	0.3058E+01	-.7259E+01
10	2.2500	0.1803E+01	-.1788E+01	-.3606E+01	0.3588E+00	-.1685E+00	-.1855E+02
11	2.5000	0.1084E+01	-.2557E+01	-.2549E+01	-.2105E+00	-.2382E+01	0.8438E+00
12	2.7500	0.3046E+00	-.2996E+01	-.9614E+00	0.5659E+00	0.4142E+00	0.2153E+02
13	3.0000	-.3400E+00	-.2847E+01	0.2152E+01	0.2052E+01	0.4524E+01	0.1135E+02
14	3.2500	-.6363E+00	-.1882E+01	0.5572E+01	0.2337E+01	0.3543E+01	-.1920E+02
15	3.5000	-.5110E+00	-.3421E+00	0.6746E+01	0.7394E+00	-.2625E+01	-.3013E+02
16	3.7500	-.6742E-01	0.1098E+01	0.4772E+01	-.1231E+01	-.7137E+01	-.5967E+01
17	4.0000	0.4318E+00	0.1886E+01	0.1531E+01	-.1764E+01	-.5007E+01	0.2301E+02
18	4.2500	0.9161E+00	0.2007E+01	-.5596E+00	-.9579E+00	0.5471E+00	0.2142E+02
19	4.5000	0.1325E+01	0.1786E+01	-.1206E+01	-.4066E+00	0.2715E+01	-.4080E+01
20	4.7500	0.1610E+01	0.1400E+01	-.1882E+01	-.8160E+00	0.2836E+00	-.1537E+02
21	5.0000	0.1710E+01	0.7696E+00	-.3165E+01	-.9849E+00	-.1157E+01	0.3850E+01
22	5.2500	0.1577E+01	-.7782E-01	-.3614E+01	0.4299E+00	0.2492E+01	0.2534E+02
23	5.5000	0.1332E+01	-.7549E+00	-.1802E+01	0.2808E+01	0.7586E+01	0.1542E+02
24	5.7500	0.1169E+01	-.8162E+00	0.1311E+01	0.4079E+01	0.7298E+01	-.1773E+02
25	6.0000	0.1198E+01	-.2684E+00	0.3071E+01	0.3241E+01	0.8649E+00	-.3374E+02

11.21

At  $x=0$  (at node 1):

$$\frac{d}{dx} \left[ EI \frac{d^2 W(x)}{dx^2} \right] = EI \frac{d^3 W}{dx^3} = -k_1 W(x)$$

$$EI \frac{d^2 W}{dx^2} = -k_{t1} \frac{dW(x)}{dx}$$

$$\text{i.e.} \quad \frac{EI}{2h^3} (W_3 - 2W_2 + 2W_{-1} - W_{-2}) = -k_1 \cdot W_1$$

$$\frac{EI}{h^2} (W_2 - 2W_1 + W_{-1}) = -k_{t1} \cdot \frac{l}{2h} (W_2 - W_{-1})$$

At  $x=l$  (at node  $n$ ):

$$\frac{d}{dx} \left[ EI \frac{d^2 W(x)}{dx^2} \right] = EI \frac{d^3 W}{dx^3} = k_2 W(x)$$

$$EI \frac{d^2 W}{dx^2} = k_{t2} \frac{dW(x)}{dx}$$

$$\text{i.e.} \quad \frac{EI}{2h^3} (W_{n+2} - 2W_{n+1} + 2W_{n-1} - W_{n-2}) = k_2 \cdot W_n$$

$$\frac{EI}{h^2} (W_{n+1} - 2W_n + W_{n-1}) = k_{t2} \cdot \frac{l}{2h} (W_{n+1} - W_{n-1})$$

11.22 Given equations can be expressed as

$$\frac{d\vec{X}}{dt} = \begin{Bmatrix} f_1 \\ f_2 \\ f_3 \\ f_4 \end{Bmatrix} = \begin{Bmatrix} x_2 \\ \frac{1}{2}(5-6x_1+2x_3) \\ x_4 \\ 20 \sin 5t + 2x_1 - 4x_3 \end{Bmatrix}$$

where

$$\vec{X} = \begin{Bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{Bmatrix} = \begin{Bmatrix} \text{original } x_1 \\ \text{original } \dot{x}_1 \\ \text{original } x_2 \\ \text{original } \dot{x}_2 \end{Bmatrix} \quad \text{and} \quad \vec{X}_0 = \begin{Bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{Bmatrix}$$

The main program which calls RK4, the subroutine FUN and the results are given.

```

C =====
C
C MAIN PROGRAM FOR CALLING THE SUBROUTINE RK4
C
C =====
C THE FOLLOWING 7 LINES NEED TO BE CHANGED FOR A DIFFERENT PROBLEM
  DIMENSION TIME(25),X(25,4),XX(4),F(4),YI(4),YJ(4),YK(4),YL(4),
2  UU(4)
  XX(1)=0.0
  XX(2)=0.0
  XX(3)=0.0
  XX(4)=0.0
  NEQ=4
  NSTEP=25
    
```

```

DT=0.25
T=0.0
WRITE (60,10)
10  FORMAT (//,3X,5H I ,10H  TIME(I),7X,5H X(1),12X,5H X(2),
2    12X,5H X(3),12X,5H X(4),/)
DO 40 I=1,NSTEP
CALL RK4 (T,DT,NEQ,XX,F,YI,YJ,YK,YL,UU)
TIME(I)=T
DO 20 J=1,NEQ
20  X(1,J)=XX(J)
WRITE (60,30) I,TIME(I),(X(1,J),J=1,NEQ)
30  FORMAT (2X,I5,F10.4,4(2X,E15.8))
40  CONTINUE
STOP
END

```

```

SUBROUTINE FUN (X,F,N,T)
DIMENSION X(N),F(N)
F(1)=X(2)
F(2)=(5.0-6.0*X(1)+2.0*X(3))/2.0
F(3)=X(4)
F(4)=20.0*SIN(5.0*T)+2.0*X(1)-4.0*X(3)
RETURN
END


```

I	TIME(I)	X(1)	X(2)	X(3)	X(4)
1	0.2500	0.76904297E-01	0.62070566E+00	0.24460433E+00	0.26932180E+01
2	0.5000	0.31453162E+00	0.13009652E+01	0.14515579E+01	0.64889636E+01
3	0.7500	0.74014312E+00	0.21077421E+01	0.29793470E+01	0.45641656E+01
4	1.0000	0.13591884E+01	0.27739763E+01	0.32607102E+01	-0.26126151E+01
5	1.2500	0.20711818E+01	0.27800636E+01	0.18916913E+01	-0.72935324E+01
6	1.5000	0.26687763E+01	0.18683126E+01	0.27935052E+00	-0.45067854E+01
7	1.7500	0.29524713E+01	0.35540020E+00	-0.78577667E-01	0.14555850E+01
8	2.0000	0.28473990E+01	-0.11584096E+01	0.60296357E+00	0.29186647E+01
9	2.2500	0.24058905E+01	-0.23112063E+01	0.90024042E+00	-0.96232796E+00
10	2.5000	0.17232550E+01	-0.30908332E+01	0.23876321E+00	-0.35336885E+01
11	2.7500	0.89089936E+00	-0.34831977E+01	-0.34061307E+00	-0.26938438E+00
12	3.0000	0.34346521E-01	-0.32377021E+01	0.25854278E+00	0.46597433E+01
13	3.2500	-0.65400219E+00	-0.21399481E+01	0.14909362E+01	0.39903281E+01
14	3.5000	-0.98161954E+00	-0.43759835E+00	0.17313157E+01	-0.24955246E+01
15	3.7500	-0.87797028E+00	0.11996664E+01	0.39478755E+00	-0.72439308E+01
16	4.0000	-0.43708453E+00	0.22119966E+01	-0.12451535E+01	-0.47495975E+01
17	4.2500	0.16884252E+00	0.25470126E+01	-0.16587874E+01	0.13506827E+01
18	4.5000	0.80284268E+00	0.24732747E+01	-0.92830122E+00	0.34720352E+01
19	4.7500	0.13844182E+01	0.21252785E+01	-0.39376926E+00	0.36745048E+00
20	5.0000	0.18346303E+01	0.14129710E+01	-0.64064902E+00	-0.15653036E+01
21	5.2500	0.20612290E+01	0.37833023E+00	-0.64773440E+00	0.23636723E+01
22	5.5000	0.20297880E+01	-0.56775379E+00	0.71066928E+00	0.81037455E+01
23	5.7500	0.18273093E+01	-0.94346517E+00	0.28840952E+01	0.79655704E+01
24	6.0000	0.16137714E+01	-0.69621664E+00	0.40854921E+01	0.10114455E+01
25	6.2500	0.14960963E+01	-0.25771955E+00	0.34342446E+01	-0.54408941E+01

11.23

Equation of motion

$$\frac{d^2 U}{dx^2} + \alpha^2 U = 0$$

At grid point  $i$ , this becomes

$$U_{i+1} - 2U_i + U_{i-1} + \alpha^2 h^2 U_i = 0 \quad \text{or} \quad U_{i+1} - (2-\lambda)U_i + U_{i-1} = 0 \quad \dots (E_1)$$

Eg. (E<sub>1</sub>), when applied to nodes 2, 3 and 4, gives where  $\lambda = \alpha^2 h^2 = \rho l^2 \omega^2 / (16E)$ .

$$\left. \begin{aligned} U_3 - (2-\lambda)U_2 + U_1 &= 0 \\ U_4 - (2-\lambda)U_3 + U_2 &= 0 \\ U_5 - (2-\lambda)U_4 + U_3 &= 0 \end{aligned} \right\} \quad \dots (E_2)$$

With boundary conditions  $U_1 = U_5 = 0$ , (E<sub>2</sub>) becomes

$$[A] \vec{U} - \lambda [I] \vec{U} = \vec{0} \quad \text{where} \quad [A] = \begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix}, \quad [I] = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad \vec{U} = \begin{Bmatrix} U_2 \\ U_3 \\ U_4 \end{Bmatrix}$$

3x3 3x1 3x3 3x1 3x1

The solution of this eigenvalue problem is found using Program 9.F. The results are



$$\begin{aligned} \lambda_1 &= 0.585786, & \lambda_2 &= 2.0, & \lambda_3 &= 3.41421 \\ \omega_1 &= 3.06147 \sqrt{\frac{E}{\rho l^2}}, & \omega_2 &= 5.65685 \sqrt{\frac{E}{\rho l^2}}, & \omega_3 &= 7.39103 \sqrt{\frac{E}{\rho l^2}} \\ \vec{U}^{(1)} &= \begin{Bmatrix} 0.5 \\ 0.707 \\ 0.5 \end{Bmatrix}, & \vec{U}^{(2)} &= \begin{Bmatrix} 0.707 \\ 0 \\ -0.707 \end{Bmatrix}, & \vec{U}^{(3)} &= \begin{Bmatrix} 0.5 \\ -0.707 \\ 0.5 \end{Bmatrix} \end{aligned}$$

**11.24** (i) Forced longitudinal vibration:

Equation is

$$EA \frac{\partial^2 u}{\partial x^2} + f = \rho A \frac{\partial^2 u}{\partial t^2} \quad \dots (E_1)$$

where  $u = u(x, t)$ .  
Let the solution be required for  $0 \leq t \leq T$ .  
Set up the finite difference grid along  $x$  and  $t$  axes as shown in Fig. 2. Note that imaginary grid points are set up at  $x = nh$  and  $t = -\Delta t$  so that free boundary conditions can be applied at  $x = (n-1)h$



and initial conditions can be applied at  $t=0$ .

Let  $u_{i,j}$  = value of  $u$  at the grid point  $(i,j)$ .

Eq. (E<sub>1</sub>) can be approximated at grid point  $(i,j)$  as

$$EA \left( \frac{u_{i+1,j} - 2u_{i,j} + u_{i-1,j}}{h^2} \right) + f_{i,j} = \rho A \left( \frac{u_{i,j+1} - 2u_{i,j} + u_{i,j-1}}{\Delta t^2} \right) \dots (E_2)$$

where  $f_{i,j} = f(x=x_i, t=t_j)$ .

The procedure to set up the finite difference equations is:

(1). Apply (E<sub>2</sub>) at grid points  $(2,1), (3,1), \dots, (n,1); (2,2), (3,2), \dots, (n,2); \dots; (2,m-1), (3,m-1), \dots, (n,m-1)$ . This gives  $(n-1 \times m-1)$  equations in the unknowns associated with all the grid points shown in Fig. 2.

(2). Apply the boundary conditions.

At  $x=0, u(x,t) = 0 \Rightarrow u_{1,1} = u_{1,2} = \dots = u_{1,m-1} = u_{1,m} = 0$

At  $x=l, \frac{\partial u}{\partial x} = 0 \Rightarrow u_{n+1,j} = u_{n-1,j}$  for  $j = 1, 2, \dots, m$

(3). Apply the initial conditions.

At  $t=0$ , let  $u(x,t) = U_0(x)$ . Then  $u_{i,1} = U_0(x=x_i)$  for  $i = 1, 2, \dots, n+1$

At  $t=0$ , let  $\frac{\partial u}{\partial t}(x,t) = \dot{U}_0(x)$ . Then  $\frac{u_{i,2} - u_{i,1}}{2 \Delta t} = \dot{U}_0(x=x_i)$

i.e.  $u_{i,-1} = u_{i,2} - 2 \Delta t \dot{U}_0(x_i)$  for  $i = 1, 2, \dots, n+1$ .

(4). Express the resulting equations in matrix form. There will be as many linear equations as there are unknowns.

(ii) Free vibration:

Equation is  $\frac{d^2 U}{dx^2} + \alpha^2 U = 0$  where  $\alpha^2 = \frac{\rho \omega^2}{E}$  --- (E<sub>3</sub>)

At node  $i$ , (E<sub>3</sub>) becomes (Fig. 1)

$U_{i+1} - 2U_i + U_{i-1} + \lambda U_i = 0$  where  $\lambda = h^2 \alpha^2$  --- (E<sub>4</sub>)

Applying (E<sub>2</sub>) at nodes 2, 3, ..., n gives

$$\left. \begin{aligned} U_3 - (2-\lambda)U_2 + U_1 &= 0 \\ U_4 - (2-\lambda)U_3 + U_2 &= 0 \\ \vdots \\ U_n - (2-\lambda)U_{n-1} + U_{n-2} &= 0 \\ U_{n+1} - (2-\lambda)U_n + U_{n-1} &= 0 \end{aligned} \right\} \dots (E_5)$$

Boundary conditions are  $U_1 = 0$  ( $U=0$  at  $x=0$ ) and  $U_{n+1} = U_{n-1}$  ( $\frac{dU}{dx} = 0$  at  $x=l$ ) --- (E<sub>6</sub>)

(E<sub>5</sub>) and (E<sub>6</sub>) give

$$\begin{aligned} U_3 - (2-\lambda)U_2 &= 0 \\ U_4 - (2-\lambda)U_3 + U_2 &= 0 \end{aligned}$$

$$\begin{aligned} & \vdots \\ U_n - (2-\lambda)U_{n-1} + U_{n-2} &= 0 \quad \dots (E_7) \\ -(2-\lambda)U_n + 2U_{n-1} &= 0 \end{aligned}$$

For  $n=4$ ,  $(E_7)$  can be expressed as

$$\begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -2 & 2 \end{bmatrix} \begin{Bmatrix} U_2 \\ U_3 \\ U_4 \end{Bmatrix} - \lambda \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{Bmatrix} U_2 \\ U_3 \\ U_4 \end{Bmatrix} = \begin{Bmatrix} 0 \\ 0 \\ 0 \end{Bmatrix} \quad \dots (E_8)$$

Frequency equation is  $\lambda^3 - 6\lambda^2 + 9\lambda - 2 = 0 \quad \dots (E_9)$

Roots are  $\lambda_1 = 0.267944$ ,  $\lambda_2 = 2.0$ ,  $\lambda_3 = 3.732050$

$$\therefore \omega_1 = 1.55289 \sqrt{\frac{E}{\rho l^2}}, \quad \omega_2 = 4.24264 \sqrt{\frac{E}{\rho l^2}}, \quad \omega_3 = 5.79555 \sqrt{\frac{E}{\rho l^2}}$$

11.25

Forced torsional vibration:

$$\text{Equation } GJ \frac{\partial^2 \theta}{\partial x^2} + f = J_0 \frac{\partial^2 \theta}{\partial t^2} \quad \dots (E_1)$$

Let the solution be required for  $0 \leq t \leq T$ .

Set up a finite difference grid along

$x$  (with  $n$  points) and along  $t$

(with  $m$  points) similar to Fig. 2 of problem 10.24.

Let  $\theta_{i,j} = \theta(x=x_i, t=t_j)$  and  $f_{i,j} = f(x=x_i, t=t_j)$

Eq.  $(E_1)$  at grid point  $(i, j)$  gives

$$GJ \left( \frac{\theta_{i+1,j} - 2\theta_{i,j} + \theta_{i-1,j}}{h^2} \right) + f_{i,j} = J_0 \left( \frac{\theta_{i,j+1} - 2\theta_{i,j} + \theta_{i,j-1}}{\Delta t^2} \right) \quad \dots (E_2)$$

The procedure to set up the finite difference grid is:

(1). Apply Eq.  $(E_2)$  at grid points  $(2,1), (3,1), \dots, (n-1,1); (2,2), (3,2), \dots, (n-1,2); \dots; (2,m-1), (3,m-1), \dots, (n-1,m-1)$ . This gives  $(n-2) \times (m-1)$  equations in the  $(n) \times (m+1)$  unknowns.

(2). Apply boundary conditions:

At  $x=0$ ,  $\theta(x,t) = 0$ . Hence  $\theta_{1,j} = 0$  for  $j = 1, 2, \dots, m$

At  $x=l$ ,  $\theta(x,t) = 0$ . Hence  $\theta_{n,j} = 0$  for  $j = 1, 2, \dots, m$

(3). Apply initial conditions.


At  $t=0$ , let  $\theta(x,t) = \theta_0(x) \Rightarrow \theta_{i,1} = \theta_0(x=x_i)$  for  $i = 1, 2, \dots, n$

At  $t=0$ , let  $\frac{\partial \theta}{\partial t}(x,t) = \dot{\theta}_0(x) \Rightarrow \theta_{i,-1} = \theta_{i,2} - 2\Delta t \dot{\theta}_0(x_i)$  for  $i = 1, 2, \dots, n$ .

(4) Express the resulting equations in matrix form. There will be  $(n-2) \times m$  equations in  $(n-2) \times m$  unknowns.

11.26

Applying Eq. (11.50) to mesh points 2, 3 and 4 gives



$$\left. \begin{aligned} w_4 - 4w_3 + (6-\lambda)w_2 - 4w_1 + w_{-1} &= 0 \\ w_5 - 4w_4 + (6-\lambda)w_3 - 4w_2 + w_1 &= 0 \\ w_6 - 4w_5 + (6-\lambda)w_4 - 4w_3 + w_2 &= 0 \end{aligned} \right\} \quad (E_1)$$

Boundary conditions are  $w = \frac{dw}{dx} = 0$  at  $x=0$  and  $x=l$

i.e.  $w_1 = 0, w_{-1} = w_2; w_5 = 0, w_6 = w_4$

Eq. (E<sub>1</sub>) becomes, after applying boundary conditions,

$$\begin{bmatrix} 7 & -4 & 1 \\ -4 & 6 & -4 \\ 1 & -4 & 7 \end{bmatrix} \begin{Bmatrix} w_2 \\ w_3 \\ w_4 \end{Bmatrix} - \lambda \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{Bmatrix} w_2 \\ w_3 \\ w_4 \end{Bmatrix} = \begin{Bmatrix} 0 \\ 0 \\ 0 \end{Bmatrix} \quad (E_2)$$

Solution (obtained using Program 9.F) is

$$\lambda_1 = 1.25544, \quad \lambda_2 = 6.0, \quad \lambda_3 = 12.7446$$

$$\omega_1 = 17.9274 \sqrt{\frac{EI}{\rho A l^4}}, \quad \omega_2 = 39.1918 \sqrt{\frac{EI}{\rho A l^4}}, \quad \omega_3 = 57.1193 \sqrt{\frac{EI}{\rho A l^4}}$$

$$\vec{w}^{(1)} = \begin{Bmatrix} 0.4544 \\ 0.7662 \\ 0.4544 \end{Bmatrix}, \quad \vec{w}^{(2)} = \begin{Bmatrix} 0.7071 \\ 0 \\ -0.7071 \end{Bmatrix}, \quad \vec{w}^{(3)} = \begin{Bmatrix} 0.5418 \\ -0.6426 \\ 0.5418 \end{Bmatrix}$$

11.27

Equation:

$$EI \frac{\partial^4 w}{\partial x^4} + \rho A \frac{\partial^2 w}{\partial t^2} = f(x, t) \quad \dots (E_1)$$



Let the solution be required for  $0 \leq t \leq T$ .

$$h = \left(\frac{l}{n-1}\right), \quad \Delta t = \frac{T}{m-1}$$

Set up a finite difference grid along

$x$  (grid points  $-1, 1, 2, \dots, n+2$ ) and along  $t$  (grid points  $-1, 1, 2, \dots, m$ ).

Imaginary grid points are located at  $x = -h, x = l+h$  and  $x = l+2h$ , and at  $t = -\Delta t$ .

Let  $w_{i,j} = w(x = x_i, t = t_j)$  and  $f_{i,j} = f(x = x_i, t = t_j)$ .

Eq. (E<sub>1</sub>) at grid point  $(i, j)$  gives

$$EI \left( \frac{w_{i+2,j} - 4w_{i+1,j} + 6w_{i,j} - 4w_{i-1,j} + w_{i-2,j}}{h^4} \right) + \rho A \left( \frac{w_{i,j+1} - 2w_{i,j} + w_{i,j-1}}{\Delta t^2} \right) = f_{i,j} = f_0 \cos \{\omega(j-1)\Delta t\} \quad \dots (E_2)$$

The procedure to set up the finite difference equations is:

(1) Apply (E<sub>2</sub>) at grid points  $(2,1), (3,1), \dots, (n,1); (2,2), (3,2), \dots, (n,2); \dots; (2,m-1), (3,m-1), \dots, (n,m-1)$ . This gives  $(n-1) \times (m-1)$  equations in the unknowns associated with all grid points.

(2). Apply boundary conditions:

$$\text{At } x=0, w = \frac{\partial w}{\partial x} = 0 \Rightarrow w_{1,j} = 0, w_{-1,j} = w_{2,j} \quad \text{for } j=1,2,\dots,m$$

$$\text{At } x=l, \frac{\partial^2 w}{\partial x^2} = \frac{\partial^3 w}{\partial x^3} = 0 \Rightarrow w_{n+1,j} - 2w_{n,j} + w_{n-1,j} = 0,$$

$$w_{n+2,j} - 2w_{n+1,j} + 2w_{n-1,j} - w_{n-2,j} = 0$$

$$\Rightarrow w_{n+1,j} = 2w_{n,j} - w_{n-1,j}, \quad w_{n+2,j} = 4w_{n,j} - 4w_{n-1,j} + w_{n-2,j}$$

for  $j=1,2,\dots,m$

(3). Apply initial conditions:

$$\text{At } t=0, \text{ let } w(x,t) = W_0(x) \Rightarrow w_{i,1} = W_0(x=x_i) \quad \text{for } i=1,2,\dots,n$$

$$\text{At } t=0, \text{ let } \frac{\partial w}{\partial t}(x,t) = \dot{W}_0(x) \Rightarrow w_{i,-1} = w_{i,2} - 2\Delta t \dot{W}_0(x_i) \quad \text{for } i=1,2,\dots,n$$

(4). Express the resulting equations in matrix form.

11.28

Equation:

$$P \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} \right) + f = \rho \frac{\partial^2 w}{\partial t^2} \quad \text{--- (E1)}$$


Set up a finite difference grid with  $n, m$  and  $p$  points along  $x, y$  and  $t$  axes. Introduce an imaginary grid point at  $t = -\Delta t$  to apply the initial conditions.

$$\text{Let } w_{i,j,k} = w(x=x_i, y=y_j, t=t_k)$$

$$\text{and } f_{i,j,k} = f(x=x_i, y=y_j, t=t_k).$$

(E1) gives at grid point  $(i,j,k)$ :

$$P \left( \frac{w_{i+1,j,k} - 2w_{i,j,k} + w_{i-1,j,k}}{h_1^2} \right) + P \left( \frac{w_{i,j+1,k} - 2w_{i,j,k} + w_{i,j-1,k}}{h_2^2} \right) + f_{i,j,k} = \rho \left( \frac{w_{i,j,k+1} - 2w_{i,j,k} + w_{i,j,k-1}}{\Delta t^2} \right) \quad \text{--- (E2)}$$



The following procedure is used to derive the final equations:

(1). Apply Eq. (E2) at grid points  $(i,j,k)$  for  $i=1,2,\dots,n$ ;  $j=1,2,\dots,m$  and  $k=1,2,\dots,p-1$ . This gives  $n \times m \times (p-1)$  equations in  $n \times m \times (p+1)$  unknowns.

(2). Apply boundary conditions:

$$\text{At } x=0, w(x,y,t) = 0 \Rightarrow w_{1,j,k} = 0 \quad \text{for } j=1,2,\dots,m; k=1,2,\dots,p$$

$$\text{At } x=a, w(x,y,t) = 0 \Rightarrow w_{n,j,k} = 0 \quad \text{for } j=1,2,\dots,m; k=1,2,\dots,p$$

At  $y=0$ ,  $w(x,y,t)=0 \Rightarrow w_{i,1,k}=0$  for  $i=1,2,\dots,n$ ;  $k=1,2,\dots,p$

At  $y=b$ ,  $w(x,y,t)=0 \Rightarrow w_{i,m,k}=0$  for  $i=1,2,\dots,n$ ;  $k=1,2,\dots,p$

(3). Apply initial conditions:

At  $t=0$ , let  $w(x,y,t)=w_0(x,y) \Rightarrow w_{i,j,1}=w_0(x_i,y_j)$  for  $i=1,2,\dots,n$ ;  
 $j=1,2,\dots,m$

At  $t=0$ , let  $\frac{\partial w}{\partial t}(x,y,t)=\dot{w}_0(x,y) \Rightarrow w_{i,j,-1}=w_{i,j,2}-2 \Delta t \dot{w}_0(x_i,y_j)$   
 for  $i=1,2,\dots,n$ ;  $j=1,2,\dots,m$

(4). Express the resulting equations in matrix form. There will be  $(n-2) \times (m-2) \times p$  equations in the same number of unknowns.

11.29 Equations of motion:

$$\begin{bmatrix} m_1 & 0 \\ 0 & m_2 \end{bmatrix} \begin{Bmatrix} \ddot{x}_1 \\ \ddot{x}_2 \end{Bmatrix} + \begin{bmatrix} c & -c \\ -c & c \end{bmatrix} \begin{Bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{Bmatrix} + \begin{bmatrix} k_1+k & -k \\ -k & k+k_2 \end{bmatrix} \begin{Bmatrix} x_1 \\ x_2 \end{Bmatrix} = \begin{Bmatrix} F_1(t) \\ F_2(t) \end{Bmatrix}$$

i.e.,

$$\begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} \ddot{\vec{x}} + \begin{bmatrix} 2 & -2 \\ -2 & 2 \end{bmatrix} \dot{\vec{x}} + \begin{bmatrix} 6 & -2 \\ -2 & 8 \end{bmatrix} \vec{x} = \begin{Bmatrix} 0 \\ 10 \end{Bmatrix} \quad (1)$$

Equations (1) can be rewritten as

$$\ddot{x}_1 + 2\dot{x}_1 - 2\dot{x}_2 + 6x_1 - 2x_2 = 0$$

$$2\ddot{x}_2 - 2\dot{x}_1 + 2\dot{x}_2 - 2x_1 + 8x_2 = 10$$

or  $\ddot{x}_1 = -2\dot{x}_1 + 2\dot{x}_2 - 6x_1 + 2x_2 \quad (2)$

$$\ddot{x}_2 = \dot{x}_1 - \dot{x}_2 + x_1 - 4x_2 + 5 \quad (3)$$


Using

$$\vec{y} = \begin{Bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{Bmatrix} = \begin{Bmatrix} x_1 \\ \dot{x}_1 \\ x_2 \\ \dot{x}_2 \end{Bmatrix} \quad \text{and} \quad \vec{Y}(t=0) = \vec{0}$$

Eqs. (2) and (3) can be expressed as

$$\frac{d\vec{Y}}{dt} = \begin{Bmatrix} y_2 \\ -2y_2 + 2y_4 - 6y_1 + 2y_3 \\ y_4 \\ y_2 - y_4 + y_1 - 4y_3 + 5 \end{Bmatrix} = \vec{f}(t)$$

Using  $n=4$ ,  $xx = \begin{Bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{Bmatrix}$ ,  $dt = \frac{\pi}{20}$ , Program 14.m can be used to find the solution.



Results of Ex11\_29

\*\*\*\*\*

>> program14

I	Time(I)	x(1)	dx(1)	x(2)	dx(2)
1	1.5708e-001	5.9523e-003	1.0944e-001	5.8328e-002	7.2086e-001
2	3.1416e-001	4.2823e-002	3.7884e-001	2.1918e-001	1.3032e+000
3	4.7124e-001	1.2857e-001	7.1703e-001	4.5992e-001	1.7366e+000
4	6.2832e-001	2.6725e-001	1.0403e+000	7.5658e-001	2.0145e+000
5	7.8540e-001	4.5102e-001	1.2817e+000	1.0845e+000	2.1343e+000
6	9.4248e-001	6.6316e-001	1.3960e+000	1.4189e+000	2.0980e+000
7	1.0996e+000	8.8165e-001	1.3607e+000	1.7358e+000	1.9137e+000
8	1.2566e+000	1.0827e+000	1.1755e+000	2.0132e+000	1.5971e+000
9	1.4137e+000	1.2440e+000	8.5854e-001	2.2318e+000	1.1714e+000
10	1.5708e+000	1.3472e+000	4.4240e-001	2.3770e+000	6.6715e-001
11	1.7279e+000	1.3799e+000	-3.0920e-002	2.4392e+000	1.2076e-001
12	1.8850e+000	1.3369e+000	-5.1500e-001	2.4149e+000	-4.2760e-001
13	2.0420e+000	1.2200e+000	-9.6373e-001	2.3069e+000	-9.3696e-001
14	2.1991e+000	1.0382e+000	-1.3356e+000	2.1246e+000	-1.3687e+000
15	2.3562e+000	8.0625e-001	-1.5969e+000	1.8828e+000	-1.6895e+000
16	2.5133e+000	5.4353e-001	-1.7248e+000	1.6009e+000	-1.8747e+000
17	2.6704e+000	2.7201e-001	-1.7082e+000	1.3017e+000	-1.9098e+000
18	2.8274e+000	1.4410e-002	-1.5489e+000	1.0090e+000	-1.7921e+000
19	2.9845e+000	-2.0779e-001	-1.2607e+000	7.4627e-001	-1.5312e+000
20	3.1416e+000	-3.7614e-001	-8.6809e-001	5.3442e-001	-1.1482e+000
21	3.2987e+000	-4.7674e-001	-4.0407e-001	3.9037e-001	-6.7391e-001
22	3.4558e+000	-5.0135e-001	9.2720e-002	3.2551e-001	-1.4669e-001
23	3.6128e+000	-4.4804e-001	5.8124e-001	3.4484e-001	3.9092e-001

```

24 3.7699e+000 -3.2129e-001 1.0214e+000 4.4658e-001 8.9549e-001
25 3.9270e+000 -1.3159e-001 1.3773e+000 6.2229e-001 1.3263e+000
26 4.0841e+000 1.0545e-001 1.6202e+000 8.5755e-001 1.6487e+000
27 4.2412e+000 3.7042e-001 1.7305e+000 1.1332e+000 1.8367e+000
28 4.3982e+000 6.4167e-001 1.6997e+000 1.4267e+000 1.8755e+000
29 4.5553e+000 8.9711e-001 1.5307e+000 1.7143e+000 1.7622e+000
30 4.7124e+000 1.1160e+000 1.2378e+000 1.9728e+000 1.5063e+000
31 4.8695e+000 1.2807e+000 8.4493e-001 2.1811e+000 1.1290e+000
32 5.0265e+000 1.3779e+000 3.8448e-001 2.3227e+000 6.6103e-001
33 5.1836e+000 1.3999e+000 -1.0601e-001 2.3860e+000 1.4069e-001
34 5.3407e+000 1.3451e+000 -5.8663e-001 2.3663e+000 -3.8975e-001
35 5.4978e+000 1.2182e+000 -1.0184e+000 2.2653e+000 -8.8723e-001
36 5.6549e+000 1.0296e+000 -1.3664e+000 2.0915e+000 -1.3115e+000
37 5.8119e+000 7.9480e-001 -1.6025e+000 1.8590e+000 -1.6283e+000
38 5.9690e+000 5.3301e-001 -1.7080e+000 1.5869e+000 -1.8122e+000
39 6.1261e+000 2.6552e-001 -1.6747e+000 1.2974e+000 -1.8486e+000
40 6.2832e+000 1.4057e-002 -1.5055e+000 1.0141e+000 -1.7349e+000
%=====
%
% Program14.m
% Main program for calling the subroutine RK4
%
%=====
% Run "Program14" in MATLAB command window. Program14.m, rk4.m, and fun.m
% should be in the same folder, and set the Matlab path to this folder
% following 5 lines contain problem-dependent data
format long
xx=[0 0 0 0];
neq=4;
nstep=40;
dt=pi/20;
t=0;
%end of problem-dependent data
fprintf(' I    Time(I)    x(1)    dx(1)    x(2)    dx(2) \n\n');
for i=1:nstep
    [xx,f,t]=rk4(t,dt,neq,xx);
    time(i)=t;
    for j=1:neq
        x(i,j)=xx(j);
    end
    fprintf('%2.0f %8.4e %8.4e %8.4e %8.4e %8.4e\n',i,time(i),x(i,1:neq));
end
plot(time', x(1:40,1));
gtext('x(1)');
hold on;
plot(time', x(1:40,3));
gtext('x(2)');
xlabel('Time');
ylabel('X');
%=====
%
% Function rk4.m
%
%=====
function [xx,f,t]=rk4(t,dt,n,xx)
[xi]=fun(xx,n,t);
for i=1:n
    uu(i)=xx(i)+.5*dt*xi(i);
end

```

```

tn=t+0.5*dt;
[xj]=fun(uu,n,tn);
for i=1:n
    uu(i)=xx(i)+.5*dt*xj(i);
end
[xk]=fun(uu,n,tn);
for i=1:n
    uu(i)=xx(i)+dt*xk(i);
end
tn=t+dt;
[xl]=fun(uu,n,tn);
for i=1:n
    f(i)=xl(i);
    xx(i)=xx(i)+(xi(i)+2*xj(i)+2*xk(i)+xl(i))*dt/6;
end
t=t+dt;

%=====
%
% Function fun.m
%
%=====
function [f]=fun(x,n,t)
f(1)=x(2);
f(2)=-2*x(2) + 2*x(4) - 6*x(1) + 2*x(3);
f(3) = x(4);
f(4) = x(2) - x(4) + x(1) - 4*x(3) + 5;

```

11.30 Equations of motion:

$$\ddot{x}_1 + 6x_1 - 2x_2 = 10 \sin 5t$$

$$2\ddot{x}_2 - 2x_1 + 8x_2 = 0$$

or

$$\ddot{x}_1 = -6x_1 + 2x_2 + 10 \sin 5t$$

$$\ddot{x}_2 = x_1 - 4x_2$$


Problem to be solved is:

$$\frac{d\vec{Y}}{dt} = \begin{Bmatrix} y_2 \\ -6y_1 + 2y_3 + 10 \sin 5t \\ y_4 \\ y_1 - 4y_3 \end{Bmatrix} = \vec{f}$$

with

$$\vec{Y} = \begin{Bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{Bmatrix} = \begin{Bmatrix} x_1 \\ \dot{x}_1 \\ x_2 \\ \dot{x}_2 \end{Bmatrix} \quad \text{and} \quad \vec{Y}(0) = \vec{0}$$

Program14.m is used to solve these equations with  
 $n = 4$ ,  $x_0 = \begin{Bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{Bmatrix}$  and  $dt = \frac{\pi}{20}$  (40 steps used).



Results of Ex11\_30

\*\*\*\*\*

>> program14

I	Time(I)	x(1)	dx(1)	x(2)	dx(2)
1	1.5708e-001	3.1474e-002	5.7845e-001	0.0000e+000	1.2360e-003
2	3.1416e-001	2.2179e-001	1.8901e+000	1.1243e-003	1.8174e-002
3	4.7124e-001	6.1079e-001	2.9264e+000	8.0316e-003	7.9050e-002
4	6.2832e-001	1.0743e+000	2.7210e+000	2.9302e-002	2.0111e-001
5	7.8540e-001	1.3821e+000	9.7131e-001	7.3678e-002	3.6666e-001
6	9.4248e-001	1.3282e+000	-1.7201e+000	1.4387e-001	5.1794e-001
7	1.0996e+000	8.5623e-001	-4.1349e+000	2.3154e-001	5.7695e-001
8	1.2566e+000	1.0705e-001	-5.0999e+000	3.1685e-001	4.8144e-001
9	1.4137e+000	-6.4233e-001	-4.1400e+000	3.7374e-001	2.1824e-001
10	1.5708e+000	-1.1164e+000	-1.7473e+000	3.7893e-001	-1.6442e-001
11	1.7279e+000	-1.1780e+000	8.9242e-001	3.2064e-001	-5.7311e-001
12	1.8850e+000	-8.8870e-001	2.5560e+000	2.0294e-001	-9.0595e-001
13	2.0420e+000	-4.5992e-001	2.6458e+000	4.3964e-002	-1.0910e+000
14	2.1991e+000	-1.2531e-001	1.4788e+000	-1.3077e-001	-1.1074e+000
15	2.3562e+000	-9.7785e-003	4.2490e-002	-2.9627e-001	-9.7979e-001
16	2.5133e+000	-7.0457e-002	-6.2931e-001	-4.3343e-001	-7.5366e-001
17	2.6704e+000	-1.4170e-001	-8.8991e-002	-5.2993e-001	-4.6650e-001
18	2.8274e+000	-5.4399e-002	1.2595e+000	-5.7769e-001	-1.3404e-001
19	2.9845e+000	2.4398e-001	2.4172e+000	-5.6974e-001	2.4201e-001
20	3.1416e+000	6.4251e-001	2.4157e+000	-4.9981e-001	6.5101e-001
21	3.2987e+000	9.2294e-001	9.3611e-001	-3.6547e-001	1.0521e+000
22	3.4558e+000	8.8747e-001	-1.4469e+000	-1.7347e-001	1.3711e+000
23	3.6128e+000	4.8295e-001	-3.5521e+000	5.6379e-002	1.5210e+000
24	3.7699e+000	-1.5268e-001	-4.2469e+000	2.9198e-001	1.4382e+000
25	3.9270e+000	-7.5178e-001	-3.0953e+000	4.9531e-001	1.1148e+000
26	4.0841e+000	-1.0537e+000	-6.2016e-001	6.3242e-001	6.0943e-001
27	4.2412e+000	-9.4005e-001	1.9747e+000	6.8277e-001	2.9645e-002

```

28 4.3982e+000 -4.9282e-001 3.4645e+000 6.4407e-001 -5.0708e-001
29 4.5553e+000 5.7387e-002 3.2671e+000 5.3052e-001 -9.1308e-001
30 4.7124e+000 4.6138e-001 1.7307e+000 3.6614e-001 -1.1528e+000
31 4.8695e+000 5.8458e-001 -1.1398e-001 1.7658e-001 -1.2380e+000
32 5.0265e+000 4.6760e-001 -1.1837e+000 -1.6504e-002 -1.2032e+000
33 5.1836e+000 2.8161e-001 -9.8207e-001 -1.9670e-001 -1.0768e+000
34 5.3407e+000 2.0842e-001 1.2869e-001 -3.5051e-001 -8.6708e-001
35 5.4978e+000 3.1891e-001 1.1786e+000 -4.6447e-001 -5.6937e-001
36 5.6549e+000 5.2369e-001 1.2108e+000 -5.2494e-001 -1.8930e-001
37 5.8119e+000 6.2652e-001 -9.9538e-002 -5.2141e-001 2.3595e-001
38 5.9690e+000 4.4921e-001 -2.2015e+000 -4.5213e-001 6.3386e-001
39 6.1261e+000 -4.6585e-002 -3.9510e+000 -3.2835e-001 9.1654e-001
40 6.2832e+000 -7.1472e-001 -4.2611e+000 -1.7399e-001 1.0159e+000

```

```

%=====
%
% Function fun.m
%
%=====
function [f]=fun(x,n,t)
f(1)=x(2);
f(2)= - 6*x(1) + 2*x(3) + 10*sin(5*t);
f(3) = x(4);
f(4) = x(1) - 4*x(3);

```

11.31 Equations of motion:

$$[m] \ddot{\vec{x}} + [c] \dot{\vec{x}} + [k] \vec{x} = \vec{F}(t)$$

$$\text{with } [m] = \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix}, [c] = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, [k] = \begin{bmatrix} 6 & -2 \\ -2 & 4 \end{bmatrix},$$

$$\vec{F}(t) = \begin{Bmatrix} 5 \\ 20 \sin 5t \end{Bmatrix}$$

solution using Program 15.m :

$$n=2, m = \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix}, c = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, k = \begin{bmatrix} 6 & -2 \\ -2 & 4 \end{bmatrix}, x_i = \begin{Bmatrix} 0 \\ 0 \end{Bmatrix}$$

$$x_{di} = \begin{Bmatrix} 0 \\ 0 \end{Bmatrix}, nstep=24, delat = 0.25$$

$$\text{In subprogram: } \vec{F} = \begin{Bmatrix} 5 \\ 20 \sin 5t \end{Bmatrix}$$

Results of Ex11\_31

\*\*\*\*\*

>> program15

Solution by central difference method

Given data:

n= 2      nstp= 24      delat=2.500000e-001



Solution:

step	time	x(i,1)	xd(i,1)	xdd(i,1)	x(i,2)	xd(i,2)
1	0.0000	0.0000e+000	0.0000e+000	2.5000e+000	0.0000e+000	0.0000e+000
2	0.2500	7.8125e-002	0.0000e+000	2.5000e+000	0.0000e+000	0.0000e+000

```

3 0.5000 2.9785e-001 5.9570e-001 2.2656e+000 1.1960e+000 2.3920e+000
1.9136e+001
4 0.7500 6.9273e-001 1.2292e+000 2.8024e+000 2.8783e+000 5.7566e+000
7.7812e+000
5 1.0000 1.2939e+000 1.9920e+000 3.3001e+000 3.2132e+000 4.0344e+000
-2.1559e+001
6 1.2500 2.0095e+000 2.6335e+000 1.8316e+000 1.7079e+000 -2.3409e+000
-2.9444e+001
7 1.5000 2.6113e+000 2.6349e+000 -1.8206e+000 -1.4739e-002 -6.4559e+0
00 -3.4760e+000
8 1.7500 2.8788e+000 1.7387e+000 -5.3487e+000 -2.3474e-001 -3.8852e+0
00 2.4042e+001
9 2.0000 2.7482e+000 2.7373e-001 -6.3712e+000 7.4471e-001 1.5189e+000
1.9191e+001
10 2.2500 2.3050e+000 -1.1476e+000 -4.9998e+000 1.2015e+000 2.8724e+00
0 -8.3629e+000
11 2.5000 1.6610e+000 -2.1743e+000 -3.2136e+000 4.3624e-001 -6.1695e-0
01 -1.9552e+001
12 2.7500 8.8908e-001 -2.8319e+000 -2.0468e+000 -3.1334e-001 -3.0296e+
000 2.5063e-001
:
:
0 2.1734e+001
20 4.7500 1.3287e+000 2.0191e+000 -9.4959e-001 -1.5950e-001 3.4773e+00
0 -4.4812e+000
21 5.0000 1.7010e+000 1.6947e+000 -1.6456e+000 -4.5222e-001 8.7311e-00
1 -1.6352e+001
22 5.2500 1.8823e+000 1.1071e+000 -3.0551e+000 -5.8471e-001 -8.5041e-0
01 2.5638e+000
23 5.5000 1.8304e+000 2.5880e-001 -3.7315e+000 7.8790e-001 2.4802e+000
2.4081e+001
24 5.7500 1.6408e+000 -4.8304e-001 -2.2032e+000 3.0664e+000 7.3022e+00
0 1.4494e+001
25 6.0000 1.4914e+000 -6.7792e-001 6.4412e-001 4.2109e+000 6.8461e+000
-1.8143e+001

```



11.32

Equations of motion:

$$[m] \ddot{\vec{x}} + [c] \dot{\vec{x}} + [k] \vec{x} = \vec{F}(t)$$

with

$$[m] = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}, [c] = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}, [k] = \begin{bmatrix} 6 & -2 \\ -2 & 8 \end{bmatrix}, \vec{F}(t) = \begin{Bmatrix} 0 \\ 10 \end{Bmatrix}$$

Solution using Program 15.m:

$$n = 2, m = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}, c = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}, k = \begin{bmatrix} 6 & -2 \\ -2 & 8 \end{bmatrix}, x_i = \begin{Bmatrix} 0 \\ 0 \end{Bmatrix},$$

$$x_{di} = \begin{Bmatrix} 0 \\ 0 \end{Bmatrix}, nstep = 50, \text{delt} = \frac{\pi}{20}.$$

$$\text{In subprogram: } \vec{F} = \begin{Bmatrix} 0 \\ 10 \end{Bmatrix}$$

Results of Ex11\_32

\*\*\*\*\*

&gt;&gt; program15


Solution by central difference method

Given data:

n= 2      nstp= 50      delt=2.500000e-001

Solution:

step i,2)	time xdd(i,2)	x(i,1)	xd(i,1)	xdd(i,1)	x(i,2)	xd(i,2)
1	0.0000	0.0000e+000	0.0000e+000	0.0000e+000	0.0000e+000	0.0000e+000
2	0.2500	3.4694e-018	6.9389e-018	5.5511e-017	1.5625e-001	0.0000e+000
3	0.5000	7.9153e-002	1.5831e-001	1.2664e+000	5.5613e-001	1.1123e+000
4	0.7500	2.8025e-001	5.6050e-001	1.9511e+000	1.0934e+000	1.8742e+000
5	1.0000	5.8633e-001	1.0144e+000	1.6798e+000	1.6506e+000	2.1889e+000
6	1.2500	9.2647e-001	1.2924e+000	5.4495e-001	2.1205e+000	2.0542e+000
7	1.5000	1.2034e+000	1.2342e+000	-1.0110e+000	2.4211e+000	1.5410e+000
8	1.7500	1.3294e+000	8.0596e-001	-2.4148e+000	2.5053e+000	7.6957e-001
9	2.0000	1.4627e+000	5.9627e-001	-3.4627e+000	2.5053e+000	7.6957e-001
10	2.2500	1.6015e+000	4.0153e-001	-4.5015e+000	2.4211e+000	1.5410e+000
11	2.5000	1.7447e+000	2.0447e-001	-5.5447e+000	1.0934e+000	1.0934e+000
12	2.7500	1.8915e+000	0.0000e+000	-6.5915e+000	0.0000e+000	0.0000e+000
13	3.0000	2.0415e+000	0.0000e+000	-7.6415e+000	0.0000e+000	0.0000e+000
14	3.2500	2.1947e+000	0.0000e+000	-8.6947e+000	0.0000e+000	0.0000e+000
15	3.5000	2.3515e+000	0.0000e+000	-9.7515e+000	0.0000e+000	0.0000e+000
16	3.7500	2.5115e+000	0.0000e+000	-1.08115e+001	0.0000e+000	0.0000e+000
17	4.0000	2.6747e+000	0.0000e+000	-1.18747e+001	0.0000e+000	0.0000e+000
18	4.2500	2.8415e+000	0.0000e+000	-1.29415e+001	0.0000e+000	0.0000e+000
19	4.5000	3.0115e+000	0.0000e+000	-1.40115e+001	0.0000e+000	0.0000e+000
20	4.7500	3.1847e+000	0.0000e+000	-1.50847e+001	0.0000e+000	0.0000e+000
21	5.0000	3.3615e+000	0.0000e+000	-1.61615e+001	0.0000e+000	0.0000e+000
22	5.2500	3.5415e+000	0.0000e+000	-1.72415e+001	0.0000e+000	0.0000e+000
23	5.5000	3.7247e+000	0.0000e+000	-1.83247e+001	0.0000e+000	0.0000e+000
24	5.7500	3.9115e+000	0.0000e+000	-1.94115e+001	0.0000e+000	0.0000e+000
25	6.0000	4.1015e+000	0.0000e+000	-2.05015e+001	0.0000e+000	0.0000e+000
26	6.2500	4.2947e+000	0.0000e+000	-2.15947e+001	0.0000e+000	0.0000e+000
27	6.5000	4.4915e+000	0.0000e+000	-2.26915e+001	0.0000e+000	0.0000e+000
28	6.7500	4.6915e+000	0.0000e+000	-2.37915e+001	0.0000e+000	0.0000e+000
29	7.0000	4.8947e+000	0.0000e+000	-2.48947e+001	0.0000e+000	0.0000e+000
30	7.2500	5.1015e+000	0.0000e+000	-2.60015e+001	0.0000e+000	0.0000e+000
31	7.5000	5.3115e+000	0.0000e+000	-2.71115e+001	0.0000e+000	0.0000e+000
32	7.7500	5.5247e+000	0.0000e+000	-2.82247e+001	0.0000e+000	0.0000e+000
33	8.0000	5.7415e+000	0.0000e+000	-2.93415e+001	0.0000e+000	0.0000e+000
34	8.2500	5.9615e+000	0.0000e+000	-3.04615e+001	0.0000e+000	0.0000e+000
35	8.5000	6.1847e+000	0.0000e+000	-3.15947e+001	0.0000e+000	0.0000e+000
36	8.7500	6.4115e+000	0.0000e+000	-3.27415e+001	0.0000e+000	0.0000e+000
37	9.0000	6.6415e+000	0.0000e+000	-3.38915e+001	0.0000e+000	0.0000e+000
38	9.2500	6.8747e+000	0.0000e+000	-3.50447e+001	0.0000e+000	0.0000e+000
39	9.5000	7.1115e+000	0.0000e+000	-3.62015e+001	0.0000e+000	0.0000e+000
40	9.7500	7.3515e+000	0.0000e+000	-3.73615e+001	0.0000e+000	0.0000e+000
41	10.0000	7.5947e+000	0.0000e+000	-3.85247e+001	0.0000e+000	0.0000e+000
42	10.2500	7.8415e+000	0.0000e+000	-3.96915e+001	0.0000e+000	0.0000e+000
43	10.5000	8.0915e+000	0.0000e+000	-4.08615e+001	0.0000e+000	0.0000e+000
44	10.7500	8.3447e+000	0.0000e+000	-4.20347e+001	0.0000e+000	0.0000e+000
45	11.0000	2.6959e-001	1.1967e+000	1.7533e+000	9.8753e-001	1.2500e+000
46	11.2500	6.5996e-001	1.4887e+000	5.8245e-001	1.4537e+000	1.7136e+000
47	11.5000	1.0064e+000	1.4736e+000	-7.0280e-001	1.8992e+000	1.8233e+000
48	11.7500	1.2380e+000	1.1560e+000	-1.8379e+000	2.2327e+000	1.5582e+000
49	12.0000	1.3076e+000	6.0249e-001	-2.5905e+000	2.3864e+000	9.7439e-001
50	12.2500	1.2018e+000	-7.2283e-002	-2.8077e+000	2.3293e+000	1.9320e-001
51	12.5000	9.4307e-001	-7.2916e-001	-2.4473e+000	2.0743e+000	-6.2413e-001
52	12.7500	-3.1680e+000	-3.1680e+000	-2.4473e+000	2.0743e+000	-6.2413e-001



11.33 Equations of motion:  $[m] \ddot{\vec{x}} + [c] \dot{\vec{x}} + [k] \vec{x} = \vec{F}(t)$   
 with  $[m] = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$ ,  $[c] = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ ,  $[k] = \begin{bmatrix} 6 & -2 \\ -2 & 8 \end{bmatrix}$ ,  
 $\vec{F}(t) = \begin{Bmatrix} 10 \sin 5t \\ 0 \end{Bmatrix}$

Solution using Program 16.m:

$n=2$ ,  $m = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$ ,  $c = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ ,  $k = \begin{bmatrix} 6 & -2 \\ -2 & 8 \end{bmatrix}$ ,  $x_i = \begin{Bmatrix} 0 \\ 0 \end{Bmatrix}$ ,  
 $x_{di} = \begin{Bmatrix} 0 \\ 0 \end{Bmatrix}$ ,  $nstep = 50$ ,  $delt = \frac{\pi}{20}$ .

In subprogram:  $\vec{F}(t) = \begin{Bmatrix} 10 \sin 5t \\ 0 \end{Bmatrix}$

Results of Ex11\_33

\*\*\*\*\*

>> program16

Solution by Hobolt method

Given data:

$n = 2$      $nstep = 50$      $delt = 1.570796e-001$

Solution:

step (i,2)	time	x(i,1)	xd(i,1)	xdd(i,1)	x(i,2)	xd(i,2)	xdd
1	0.0000	0.0000e+000	0.0000e+000	0.0000e+000	0.0000e+000	0.0000e+000	0.0000e+000
2	0.1571	0.0000e+000	0.0000e+000	0.0000e+000	6.1685e-002	2.2087e-017	5.0000e+000

```



3 0.3142 3.0440e-003 9.6895e-003 1.2337e-001 2.4065e-001 7.6602e-001
4.7533e+000
4 0.4712 1.8912e-002 1.6259e-001 9.1608e-001 5.1478e-001 2.0011e+000
2.9598e+000
5 0.6283 5.8032e-002 3.4519e-001 1.3650e+000 8.5661e-001 2.3334e+000
1.6316e+000
6 0.7854 1.2964e-001 5.7894e-001 1.6914e+000 1.2346e+000 2.4551e+000
1.9106e-001
7 0.9425 2.3963e-001 8.3483e-001 1.7936e+000 1.6157e+000 2.3650e+000
-1.2232e+000

```

```

45 6.9115 -3.7289e-001 1.7185e-001 3.0312e+000 3.9690e-001 -1.0620e+00
0 3.0395e+000
46 7.0686 -3.1591e-001 5.6154e-001 2.4307e+000 2.6762e-001 -5.5794e-00
1 3.6136e+000
47 7.2257 -2.0460e-001 8.5380e-001 1.6722e+000 2.2229e-001 5.0005e-003
3.9062e+000
48 7.3827 -5.5446e-002 1.0351e+000 8.6621e-001 2.6677e-001 5.8147e-001
3.8775e+000
49 7.5398 1.1405e-001 1.1066e+000 1.1476e-001 3.9952e-001 1.1229e+000
3.5160e+000
50 7.6969 2.8752e-001 1.0823e+000 -5.0214e-001 6.1148e-001 1.5822e+000
2.8416e+000
51 7.8540 4.5143e-001 9.8440e-001 -9.3551e-001 8.8654e-001 1.9178e+000
1.9053e+000

```





11.34 Equations of motion: Given in solution of Problem 11.20

Solution using Program 16.m:

$$n=2, m = \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix}, c = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, k = \begin{bmatrix} 6 & -2 \\ -2 & 4 \end{bmatrix}, x_i = \begin{Bmatrix} 0 \\ 0 \end{Bmatrix},$$

$$x_{di} = \begin{Bmatrix} 0 \\ 0 \end{Bmatrix}, nstep = 50, \text{delt} = 0.25$$

$$\text{In subprogram: } \vec{F} = \begin{Bmatrix} 5 \\ 20 \sin 5t \end{Bmatrix}$$



Results of Ex11\_34

\*\*\*\*\*

>> program16

Solution by Hobolt method

Given data:

n= 2 nstp= 50 delt=2.500000e-001

Solution:

step	time	x(i,1)	xd(i,1)	xdd(i,1)	x(i,2)	xd(i,2)	xdd
1	0.0000	0.0000e+000	0.0000e+000	2.5000e+000	0.0000e+000	0.0000e+000	0.0000e+000
2	0.2500	7.8125e-002	0.0000e+000	2.5000e+000	0.0000e+000	0.0000e+000	0.0000e+000
3	0.5000	2.9785e-001	5.9570e-001	2.2656e+000	1.1960e+000	2.3920e+000	1.9136e+001
4	0.7500	6.7731e-001	1.8615e+000	2.8459e+000	2.3779e+000	3.0857e+000	-1.9588e+001
5	1.0000	1.1875e+000	2.2638e+000	1.6286e+000	2.6912e+000	-1.6232e+000	-2.7568e+001

```

.
.
.
45 11.0000 2.3544e+000 9.1504e-001 -2.2782e+000 2.2850e+000 -3.3050e+0
00 -2.4426e+001
46 11.2500 2.4907e+000 1.5041e-001 -3.4958e+000 1.4763e+000 -4.2224e+0
00 -6.8088e+000
47 11.5000 2.4280e+000 -7.0701e-001 -3.8836e+000 9.0044e-001 -6.8674e-
001 1.7538e+001
48 11.7500 2.1705e+000 -1.4140e+000 -3.0502e+000 9.6130e-001 2.0554e+0
00 1.6650e+001
49 12.0000 1.7634e+000 -1.8673e+000 -1.6701e+000 1.1202e+000 1.1342e-0
01 -7.0501e+000
50 12.2500 1.2630e+000 -2.1135e+000 -5.9375e-001 6.9518e-001 -3.7771e+
000 -2.0254e+001
51 12.5000 7.2135e-001 -2.1793e+000 1.7567e-001 -1.6028e-001 -4.0782e+
000 -4.4321e+000

```

11.35 The problem-dependent data to be used in the main program that calls WILSON (given in Problem 11.49), subroutine EXTFUN and output are given.

~~C FOLLOWING 10 LINES CONTAIN PROBLEM-DEPENDENT DATA~~

~~REAL M(2,2),MI(2,2),K(2,2)~~

~~DIMENSION C(2,2),XI(2),XDI(2),XDDI(2),X(25,2),XD(25,2),XDD(25,2),~~

~~2 XT(2),F(2),F1(2),F2(2),F3(2),R(2),LA(2),LB(2,2),S(2),ZA(2),~~

~~3 RK(2,2),XN1(2),XN2(2),XN3(2),XN4(2)~~

~~DATA N,NSTEP,NSTEP1,TH,DELTA/2,24,25,1.4,0.24216267/~~

~~DATA XI/0.0,0.0/~~

~~DATA XDI/0.0,0.0/~~

~~DATA M/1.0,0.0,0.0,2.0/~~

~~DATA C/2.0,-2.0,-2.0,2.0/~~

~~DATA K/6.0,-2.0,-2.0,8.0/~~

~~C END OF PROBLEM-DEPENDENT DATA~~

~~SUBROUTINE EXTFUN (F,TIME,N)~~

~~DIMENSION F(N)~~

~~F(1)=0.0~~

~~F(2)=10.0~~

~~RETURN~~

~~END~~

~~SOLUTION BY WILSON METHOD~~

GIVEN DATA:

~~N = 2    NSTEP = 24    TH = 0.14000000E+01    DELTA = 0.24216267E+00~~

SOLUTION:

STEP	TIME	X(1,1)	XD(1,1)	XDD(1,1)	X(1,2)	XD(1,2)	XDD(1,2)
1	0.0000	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.5000E+01
2	0.2422	0.1551E-01	0.1922E+00	0.1587E+01	0.1330E+00	0.1042E+01	0.3610E+01
3	0.4843	0.1136E+00	0.6390E+00	0.2103E+01	0.4767E+00	0.1735E+01	0.2112E+01
4	0.7265	0.3250E+00	0.1087E+01	0.1593E+01	0.9440E+00	0.2063E+01	0.5986E+00
5	0.9687	0.6233E+00	0.1328E+01	0.4052E+00	0.1447E+01	0.2036E+01	-0.8224E+00
6	1.2108	0.9431E+00	0.1256E+01	-0.9998E+00	0.1905E+01	0.1691E+01	-0.2029E+01
7	1.4530	0.1206E+01	0.8697E+00	-0.2195E+01	0.2246E+01	0.1095E+01	-0.2892E+01
8	1.6951	0.1346E+01	0.2556E+00	-0.2877E+01	0.2423E+01	0.3457E+00	-0.3297E+01
9	1.9373	0.1323E+01	-0.4452E+00	-0.2910E+01	0.2411E+01	-0.4384E+00	-0.3179E+01
10	2.1795	0.1136E+01	-0.1078E+01	-0.2318E+01	0.2218E+01	-0.1131E+01	-0.2545E+01

```

11  2.4216  0.8171E+00  -.1511E+01  -.1252E+01  0.1879E+01  -.1620E+01  -.1489E+01
12  2.6638  0.4274E+00  -.1656E+01  0.5148E-01  0.1456E+01  -.1822E+01  -.1836E+00
13  2.9060  0.4036E-01  -.1489E+01  0.1330E+01  0.1023E+01  -.1705E+01  0.1149E+01
14  3.1481  -.2713E+00  -.1044E+01  0.2341E+01  0.6543E+00  -.1291E+01  0.2276E+01
15  3.3903  -.4500E+00  -.4093E+00  0.2903E+01  0.4155E+00  -.6520E+00  0.2998E+01
16  3.6324  -.4638E+00  0.2960E+00  0.2923E+01  0.3474E+00  0.9684E-01  0.3186E+01
17  3.8746  -.3114E+00  0.9418E+00  0.2410E+01  0.4606E+00  0.8226E+00  0.2808E+01
18  4.1168  -.2189E-01  0.1411E+01  0.1468E+01  0.7335E+00  0.1396E+01  0.1931E+01
19  4.3589  0.3512E+00  0.1622E+01  0.2714E+00  0.1116E+01  0.1717E+01  0.7159E+00
20  4.6011  0.7399E+00  0.1539E+01  -.9597E+00  0.1540E+01  0.1729E+01  -.6189E+00
21  4.8433  0.1074E+01  0.1180E+01  -.2005E+01  0.1929E+01  0.1432E+01  -.1834E+01
22  5.0854  0.1294E+01  0.6124E+00  -.2680E+01  0.2213E+01  0.8814E+00  -.2711E+01
23  5.3276  0.1362E+01  -.5939E-01  -.2868E+01  0.2343E+01  0.1781E+00  -.3097E+01
24  5.5697  0.1267E+01  -.7143E+00  -.2541E+01  0.2297E+01  -.5512E+00  -.2926E+01
25  5.8119  0.1027E+01  -.1235E+01  -.1763E+01  0.2085E+01  -.1176E+01  -.2233E+01

```

11.36 The problem-dependent data to be used in the main program that calls WILSON (given in Problem 11.49), subroutine EXTFUN, and the results are given.

C FOLLOWING 10 LINES CONTAIN PROBLEM-DEPENDENT DATA

```
REAL M(2,2),MI(2,2),K(2,2)
```

```
DIMENSION C(2,2),XI(2),XD(2),XDD(2),X(25,2),XD(25,2),XDD(25,2),
```

```
2 XT(2),F(2),F1(2),F2(2),FT(2),R(2),LA(2),LB(2,2),S(2),ZA(2),
```

```
3 IN(2,2),XN1(2),XN2(2),XN3(2),XN+(2)
```

```
DATA N,NSTEP,NSTEP1,TH,DELTA/2,24,25,1.4,0.24216267/
```

```
DATA X1/0.0,0.0/
```

```
DATA XD1/0.0,0.0/
```

```
DATA M/1.0,0.0,0.0,2.0/
```

```
DATA C/0.0,0.0,0.0,0.0/
```

```
DATA K/6.0,-2.0,-2.0,8.0/
```

C END OF PROBLEM-DEPENDENT DATA

```
SUBROUTINE EXTFUN (F,TIME,N)
```

```
DIMENSION F(N)
```

```
F(1)=10.0*SIN(5.0*TIME)
```

```
F(2)=0.0
```

```
RETURN
```

```
END
```

SOLUTION BY WILSON METHOD

GIVEN DATA:

```
N= 2      NSTEP= 24      TH= 0.1400000E+01      DELTA= 0.24216267E+00
```

SOLUTION:

STEP	TIME	X(1,1)	XD(1,1)	XDD(1,1)	X(1,2)	XD(1,2)	XDD(1,2)
1	0.0000	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00
2	0.2422	0.8209E-01	0.1017E+01	0.8399E+01	0.1461E-02	0.1810E-01	0.1495E+00
3	0.4843	0.5226E+00	0.2406E+01	0.3071E+01	0.1381E-01	0.9867E-01	0.5160E+00
4	0.7265	0.1062E+01	0.1503E+01	-.1053E+02	0.5495E-01	0.2498E+00	0.7325E+00
5	0.9687	0.1074E+01	-.1584E+01	-.1496E+02	0.1330E+00	0.3781E+00	0.3265E+00
6	1.2108	0.3655E+00	-.3797E+01	-.3306E+01	0.2246E+00	0.3401E+00	-.6398E+00
7	1.4530	-.5063E+00	-.2807E+01	0.1148E+02	0.2799E+00	0.8172E-01	-.1494E+01
8	1.6951	-.8380E+00	0.1135E+00	0.1263E+02	0.2548E+00	-.2930E+00	-.1600E+01
9	1.9373	-.5584E+00	0.1706E+01	0.5215E+00	0.1431E+00	-.6043E+00	-.9714E+00
10	2.1795	-.2170E+00	0.7537E+00	-.8390E+01	-.2359E-01	-.7389E+00	-.1402E+00
11	2.4216	-.2285E+00	-.6334E+00	-.3066E+01	-.2004E+00	-.6951E+00	0.5020E+00
12	2.6638	-.3632E+00	-.3125E-01	0.8040E+01	-.3488E+00	-.5089E+00	0.1036E+01

13	2.9060	-.1304E+00	0.1973E+01	0.8514E+01	-.4357E+00	-.1847E+00	0.1641E+01
14	3.1481	0.4742E+00	0.2513E+01	-.4054E+01	-.4280E+00	0.2662E+00	0.2083E+01
15	3.3903	0.8633E+00	0.2840E+00	-.1436E+02	-.3050E+00	0.7389E+00	0.1822E+01
16	3.6324	0.5643E+00	-.2534E+01	-.8916E+01	-.8396E-01	0.1040E+01	0.6659E+00
17	3.8746	-.1616E+00	-.2845E+01	0.6349E+01	0.1726E+00	0.1017E+01	-.8543E+00
18	4.1168	-.5959E+00	-.4607E+00	0.1334E+02	0.3836E+00	0.6825E+00	-.1911E+01
19	4.3589	-.4018E+00	0.1711E+01	0.4600E+01	0.4907E+00	0.1932E+00	-.2130E+01
20	4.6011	0.2833E-01	0.1349E+01	-.7598E+01	0.4783E+00	-.2815E+00	-.1791E+01
21	4.8433	0.1332E+00	-.4775E+00	-.7484E+01	0.3623E+00	-.6571E+00	-.1311E+01
22	5.0854	-.9143E-01	-.9221E+00	0.3812E+01	0.1706E+00	-.9025E+00	-.7150E+00
23	5.3276	-.1416E+00	0.7616E+00	0.1009E+02	-.5983E-01	-.9628E+00	0.2168E+00
24	5.5697	0.2564E+00	0.2184E+01	0.1656E+01	-.2753E+00	-.7698E+00	0.1377E+01
25	5.8119	0.7047E+00	0.9850E+00	-.1156E+02	-.4136E+00	-.3408E+00	0.2166E+01

11.37 The problem-dependent data to be used in the main program that calls WILSON (given in Problem 11.49), subroutine EXIFUN, and output are given.

C FOLLOWING 10 LINES CONTAIN PROBLEM-DEPENDENT DATA

REAL M(2,2),MI(2,2),K(2,2)

DIMENSION C(2,2),XI(2),XDI(2),XDDI(2),X(25,2),XD(25,2),XDD(25,2),

2 XI(2),F(2),F1(2),F2(2),FT(2),R(2),LA(2),LB(2,2),S(2),ZA(2),

3 TK(2,2),XN1(2),XN2(2),XN3(2),XN4(2)

DATA N,NSTEP,NSTEP1,TH,DELTA/2,24,25,1.4,0.25/

DATA XI/0.0,0.0/

DATA XDI/0.0,0.0/

DATA M/2.0,0.0,0.0,1.0/

DATA C/0.0,0.0,0.0,0.0/

DATA K/6.0,-2.0,-2.0,4.0/

C END OF PROBLEM-DEPENDENT DATA

SUBROUTINE EXIFUN (F,TIME,N)

DIMENSION F(N)

F(1)=5.0

F(2)=20.0\*SIN(5.0\*TIME)

RETURN

END

SOLUTION BY WILSON METHOD

GIVEN DATA:

N= 2      NSTEP= 24      TH= 0.14000000E+01      DELTA= 0.25000000E+00

SOLUTION:

STEP	TIME	X(1,1)	XD(1,1)	XDD(1,1)	X(1,2)	XD(1,2)	XDD(1,2)
1	0.0000	0.0000E+00	0.0000E+00	0.2500E+01	0.0000E+00	0.0000E+00	0.0000E+00
2	0.2500	0.7846E-01	0.6290E+00	0.2532E+01	0.1849E+00	0.2219E+01	0.1775E+02
3	0.5000	0.3173E+00	0.1292E+01	0.2772E+01	0.1185E+01	0.5347E+01	0.7278E+01
4	0.7500	0.7246E+00	0.1956E+01	0.2539E+01	0.2473E+01	0.3847E+01	-.1928E+02
5	1.0000	0.1277E+01	0.2398E+01	0.9958E+00	0.2760E+01	-.1840E+01	-.2622E+02
6	1.2500	0.1880E+01	0.2322E+01	-.1598E+01	0.1722E+01	-.5495E+01	-.3023E+01
7	1.5000	0.2387E+01	0.1638E+01	-.3881E+01	0.5026E+00	-.3266E+01	0.2085E+02
8	1.7500	0.2666E+01	0.5579E+00	-.4757E+01	0.2826E+00	0.1285E+01	0.1556E+02
9	2.0000	0.2661E+01	-.5871E+00	-.4402E+01	0.8331E+00	0.2090E+01	-.9119E+01
10	2.2500	0.2384E+01	-.1595E+01	-.3661E+01	0.9796E+00	-.1282E+01	-.1786E+02
11	2.5000	0.1980E+01	-.2397E+01	-.2755E+01	0.3022E+00	-.3333E+01	0.1457E+01
12	2.7500	0.1212E+01	-.2882E+01	-.1122E+01	-.2795E+00	-.4970E+00	0.2123E+02
13	3.0000	0.4833E+00	-.2841E+01	0.1444E+01	0.1675E+00	0.3704E+01	0.1238E+02
14	3.2500	-.1555E+00	-.2164E+01	0.3975E+01	0.1194E+01	0.3364E+01	-.1510E+02
15	3.5000	-.5604E+00	-.1028E+01	0.5115E+01	0.1459E+01	-.1662E+01	-.2510E+02
16	3.7500	-.6636E+00	0.1787E+00	0.4536E+01	0.4757E+00	-.5336E+01	-.4289E+01

17	4.0000	-.4914E+00	0.1142E+01	0.3169E+01	-.7410E+00	-.3392E+01	0.1984E+02
18	4.2500	-.1194E+00	0.1784E+01	0.1965E+01	-.1005E+01	0.1134E+01	0.1637E+02
19	4.5000	0.3765E+00	0.2138E+01	0.8735E+00	-.4563E+00	0.2272E+01	-.7272E+01
20	4.7500	0.9225E+00	0.2166E+01	-.6563E+00	-.2094E+00	-.6720E+00	-.1628E+02
21	5.0000	0.1425E+01	0.1778E+01	-.2441E+01	-.6840E+00	-.2317E+01	0.3120E+01
22	5.2500	0.1783E+01	0.1046E+01	-.3417E+01	-.9452E+00	0.1110E+01	0.2429E+02
23	5.5000	0.1943E+01	0.2598E+00	-.2874E+01	0.1076E-01	0.6216E+01	0.1655E+02
24	5.7500	0.1934E+01	-.2762E+00	-.1414E+01	0.1780E+01	0.6725E+01	-.1248E+02
25	6.0000	0.1832E+01	-.4927E+00	-.3190E+00	0.2923E+01	0.1835E+01	-.2663E+02

11.38 The problem-dependent data to be used in the main program that calls NUMARK (given in Problem 11.50), subroutine EXTFUN, and output are given:

```

C FOLLOWING 10 LINES CONTAIN PROBLEM-DEPENDENT DATA-----
  REAL M(2,2),MI(2,2),K(2,2)
  DIMENSION C(2,2),XI(2),XDI(2),XDDI(2),X(25,2),XD(25,2),XDD(25,2),
  2 XT(2),F(2),R(2),LA(2),LB(2,2),S(2),ZA(2),
  3 TK(2,2),XN1(2),XN2(2),XN3(2),XN4(2)
  DATA N,NSTEP,NSTEP1,ALPHA,BETA,DELTA/2,24,25,0.16666667,0.5,
  2 0.24216267/
  DATA X1/0.0,0.0/
  DATA XDI/0.0,0.0/
  DATA M/1.0,0.0,0.0,2.0/
  DATA C/2.0,-2.0,-2.0,2.0/
  DATA K/6.0,-2.0,-2.0,8.0/
C END OF PROBLEM-DEPENDENT DATA---
SUBROUTINE EXTFUN (F,TIME,N)
DIMENSION F(N)
F(1)=0.0
F(2)=10.0
RETURN
END

```

## SOLUTION BY NEWMARK METHOD

## GIVEN DATA:

N= 2 NSIEP= 24 ALPHA= 0.16666667E+00 BETA= 0.50000000E+00  
 DELTA= 0.24216267E+00

## SOLUTION:

STEP	TIME	X(1,1)	XD(1,1)	XDD(1,1)	X(1,2)	XD(1,2)	XDD(1,2)
1	0.0000	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.5000E+01
2	0.2422	0.1776E-01	0.2200E+00	0.1817E+01	0.1335E+00	0.1048E+01	0.3656E+01
3	0.4843	0.1287E+00	0.7143E+00	0.2266E+01	0.4799E+00	0.1753E+01	0.2170E+01
4	0.7265	0.3613E+00	0.1178E+01	0.1568E+01	0.9532E+00	0.2093E+01	0.6342E+00
5	0.9687	0.6793E+00	0.1393E+01	0.2007E+00	0.1464E+01	0.2067E+01	-0.8511E+00
6	1.2108	0.1008E+01	0.1259E+01	-0.1303E+01	0.1927E+01	0.1704E+01	-0.2145E+01
7	1.4530	0.1263E+01	0.7991E+00	-0.2497E+01	0.2268E+01	0.1071E+01	-0.3080E+01
8	1.6951	0.1377E+01	0.1221E+00	-0.3094E+01	0.2433E+01	0.2741E+00	-0.3505E+01
9	1.9373	0.1317E+01	-0.6142E+00	-0.2987E+01	0.2398E+01	-0.5540E+00	-0.3334E+01
10	2.1795	0.1088E+01	-0.1246E+01	-0.2231E+01	0.2173E+01	-0.1270E+01	-0.2581E+01
11	2.4216	0.7330E+00	-0.1639E+01	-0.1013E+01	0.1802E+01	-0.1748E+01	-0.1365E+01
12	2.6638	0.3203E+00	-0.1712E+01	0.4055E+00	0.1353E+01	-0.1901E+01	0.9807E-01
13	2.9060	-0.6943E-01	-0.1453E+01	0.1736E+01	0.9094E+00	-0.1703E+01	0.1543E+01
14	3.1481	-0.3607E+00	-0.9133E+00	0.2721E+01	0.5536E+00	-0.1189E+01	0.2700E+01
15	3.3903	-0.4977E+00	-0.1994E+00	0.3176E+01	0.3513E+00	-0.4559E+00	0.3354E+01
16	3.6324	-0.4544E+00	0.5510E+00	0.3022E+01	0.3395E+00	0.3593E+00	0.3379E+01
17	3.8746	-0.2394E+00	0.1195E+01	0.2295E+01	0.5196E+00	0.1104E+01	0.2773E+01
18	4.1168	0.1059E+00	0.1610E+01	0.1138E+01	0.8573E+00	0.1639E+01	0.1647E+01
19	4.3589	0.5158E+00	0.1720E+01	-0.2284E+00	0.1289E+01	0.1865E+01	0.2165E+00
20	4.6011	0.9129E+00	0.1506E+01	-0.1545E+01	0.1732E+01	0.1740E+01	-0.1251E+01
21	4.8433	0.1222E+01	0.1008E+01	-0.2564E+01	0.2105E+01	0.1288E+01	-0.2478E+01
22	5.0854	0.1386E+01	0.3231E+00	-0.3096E+01	0.2337E+01	0.5967E+00	-0.3235E+01
23	5.3276	0.1374E+01	-0.4202E+00	-0.3043E+01	0.2385E+01	-0.2046E+00	-0.3382E+01
24	5.5697	0.1189E+01	-0.1081E+01	-0.2419E+01	0.2241E+01	-0.9643E+00	-0.2893E+01
25	5.8119	0.8670E+00	-0.1537E+01	-0.1342E+01	0.1933E+01	-0.1540E+01	-0.1862E+01

(11.39) The problem-dependent data to be used in the main program that calls NUMARK (given in Problem 11.50), the subroutine EXTFUN, and the output are given here:

```

C FOLLOWING 10 LINES CONTAIN PROBLEM-DEPENDENT DATA
  REAL M(2,2),HI(2,2),K(2,2)
  DIMENSION C(2,2),XI(2),XDI(2),XDDI(2),X(25,2),XD(25,2),XDD(25,2),
  2 XI(2),F(2),R(2),LA(2),LB(2,2),S(2),ZA(2),
  3 IK(2,2),XN1(2),XN2(2),XN3(2),XN4(2)
  DATA N,NSIEP,NSTEP1,ALPHA,BETA,DELTA/2,24,25,0.16666667,0.5,
  2 0.24216267/
  DATA XI/0.0,0.0/
  DATA XDI/0.0,0.0/
  DATA M/1.0,0.0,0.0,2.0/
  DATA C/0.0,0.0,0.0,0.0/
  DATA K/6.0,-2.0,-2.0,8.0/
C END OF PROBLEM-DEPENDENT DATA
SUBROUTINE EXTFUN (F,TIME,N)
  DIMENSION F(N)
  F(1)=10.0*SIN(5.0*TIME)
  F(2)=0.0
  RETURN
END

```

## SOLUTION BY NEWMARK METHOD

## GIVEN DATA:

N= 2      NSTEP= 24      ALPHA= 0.16666667E+00      BETA= 0.50000000E+00  
 DELTA= 0.24216267E+00

## SOLUTION:

STEP	TIME	X(I,1)	XD(I,1)	XDD(I,1)	X(I,2)	XD(I,2)	XDD(I,2)
1	0.0000	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00
2	0.2422	0.8642E-01	0.1071E+01	0.8842E+01	0.8129E-03	0.1007E-01	0.8317E-01
3	0.4843	0.5509E+00	0.2542E+01	0.3308E+01	0.9875E-02	0.8206E-01	0.5114E+00
4	0.7265	0.1120E+01	0.1570E+01	-0.1134E+02	0.4878E-01	0.2560E+00	0.9251E+00
5	0.9687	0.1119E+01	-0.1784E+01	-0.1636E+02	0.1345E+00	0.4383E+00	0.5807E+00
6	1.2108	0.3303E+00	-0.4220E+01	-0.3762E+01	0.2457E+00	0.4296E+00	-0.6523E+00
7	1.4530	-0.6401E+00	-0.3127E+01	0.1279E+02	0.3182E+00	0.1190E+00	-0.1913E+01
8	1.6951	-0.1003E+01	0.2052E+00	0.1473E+02	0.2886E+00	-0.3739E+00	-0.2158E+01
9	1.9373	-0.6501E+00	0.2182E+01	0.1601E+01	0.1439E+00	-0.7835E+00	-0.1226E+01
10	2.1795	-0.1786E+00	0.1284E+01	-0.9018E+01	-0.6887E-01	-0.9202E+00	0.9693E-01
11	2.4216	-0.8747E-01	-0.3477E+00	-0.4458E+01	-0.2797E+00	-0.7836E+00	0.1031E+01
12	2.6638	-0.1894E+00	-0.2747E-01	0.7103E+01	-0.4342E+00	-0.4713E+00	0.1547E+01
13	2.9060	0.2207E-01	0.1815E+01	0.8110E+01	-0.4984E+00	-0.3990E-01	0.2016E+01
14	3.1481	0.5744E+00	0.2232E+01	-0.4664E+01	-0.4456E+00	0.4895E+00	0.2357E+01
15	3.3903	0.8748E+00	-0.1784E+00	-0.1524E+02	-0.2622E+00	0.1008E+01	0.1924E+01
16	3.6324	0.4459E+00	-0.3111E+01	-0.8975E+01	0.2292E-01	0.1284E+01	0.3542E+00
17	3.8746	-0.4039E+00	-0.3220E+01	0.8070E+01	0.3241E+00	0.1121E+01	-0.1700E+01
18	4.1168	-0.8682E+00	-0.2884E+00	0.1614E+02	0.5329E+00	0.5516E+00	-0.3000E+01
19	4.3589	-0.5593E+00	0.2449E+01	0.6467E+01	0.5797E+00	-0.1601E+00	-0.2878E+01
20	4.6011	0.8159E-01	0.2258E+01	-0.8046E+01	0.4672E+00	-0.7250E+00	-0.1787E+01
21	4.8433	0.3764E+00	0.1102E+00	-0.9692E+01	0.2506E+00	-0.1017E+01	-0.6259E+00
22	5.0854	0.2284E+00	-0.8793E+00	0.1519E+01	-0.5534E-02	-0.1063E+01	0.2506E+00
23	5.3276	0.1303E+00	0.3584E+00	0.8703E+01	-0.2470E+00	-0.8969E+00	0.1118E+01
24	5.5697	0.3961E+00	0.1522E+01	0.9086E+00	-0.4220E+00	-0.5091E+00	0.2084E+01
25	5.8119	0.6650E+00	0.1770E+00	-0.1202E+02	-0.4793E+00	0.5587E-01	0.2582E+01

11.40 The problem-dependent data to be used in the main program given in Problem 11.50 (which calls NUMARK), subroutine EXTFUN and results are given.

THE FOLLOWING 10 LINES CONTAIN PROBLEM-DEPENDENT DATA

REAL M(2,2),MI(2,2),K(2,2)

DIMENSION C(2,2),XI(2),XD1(2),XDDI(2),X(25,2),XD(25,2),XDD(25,2),

2 XT(2),F(2),R(2),LA(2),LB(2,2),S(2),ZA(2),

3 TK(2,2),XN1(2),XN2(2),XN3(2),XN4(2)

DATA N,NSTEP,NSTEP1,ALPHA,BETA,DELTA/

2 2,24,25,0.16666667,0.5,0.25/

DATA XI/0.0,0.0/

DATA XD1/0.0,0.0/

DATA M/2.0,0.0,0.0,1.0/

DATA C/0.0,0.0,0.0,0.0/

DATA K/6.0,-2.0,-2.0,4.0/

C END OF PROBLEM-DEPENDENT DATA

SUBROUTINE EXTFUN (F,TIME,N)

DIMENSION F(N)

F(1)=5.0

F(2)=20.0\*SIN(5.0\*TIME)

RETURN

END

## SOLUTION BY NEWMARK METHOD

GIVEN DATA:

N= 2      NSTEP= 24      ALPHA= 0.16666667E+00      BETA= 0.50000000E+00  
 DELTA= 0.24216267E+00

SOLUTION:

STEP	TIME	X(I,1)	XD(I,1)	XDD(I,1)	X(I,2)	XD(I,2)	XDD(I,2)
1	0.0000	0.0000E+00	0.0000E+00	0.2500E+01	0.0000E+00	0.0000E+00	0.0000E+00
2	0.2500	0.7769E-01	0.6198E+00	0.2458E+01	0.1914E+00	0.2296E+01	0.1837E+02
3	0.5000	0.3129E+00	0.1276E+01	0.2789E+01	0.1228E+01	0.5553E+01	0.7683E+01
4	0.7500	0.7202E+00	0.1988E+01	0.2905E+01	0.2565E+01	0.3982E+01	-.2025E+02
5	1.0000	0.1293E+01	0.2534E+01	0.1469E+01	0.2847E+01	-.2048E+01	-.2798E+02
6	1.2500	0.1940E+01	0.2517E+01	-.1606E+01	0.1715E+01	-.6001E+01	-.3641E+01
7	1.5000	0.2488E+01	0.1742E+01	-.4594E+01	0.3703E+00	-.3674E+01	0.2226E+02
8	1.7500	0.2769E+01	0.4543E+00	-.5706E+01	0.9920E-01	0.1312E+01	0.1763E+02
9	2.0000	0.2712E+01	-.8750E+00	-.4927E+01	0.7083E+00	0.2480E+01	-.8290E+01
10	2.2500	0.2353E+01	-.1940E+01	-.3596E+01	0.9630E+00	-.8686E+00	-.1850E+02
11	2.5000	0.1768E+01	-.2694E+01	-.2435E+01	0.3681E+00	-.3089E+01	0.7367E+00
12	2.7500	0.1035E+01	-.3095E+01	-.7734E+00	-.1674E+00	-.3396E+00	0.2126E+02
13	3.0000	0.2664E+00	-.2940E+01	0.2019E+01	0.3183E+00	0.3851E+01	0.1227E+02
14	3.2500	-.3745E+00	-.2064E+01	0.4988E+01	0.1364E+01	0.3319E+01	-.1652E+02
15	3.5000	-.7216E+00	-.6614E+00	0.6231E+01	0.1566E+01	-.2150E+01	-.2722E+02
16	3.7500	-.7048E+00	0.7455E+00	0.5024E+01	0.4094E+00	-.6182E+01	-.5035E+01
17	4.0000	-.3862E+00	0.1704E+01	0.2642E+01	-.1016E+01	-.4117E+01	0.2155E+02
18	4.2500	0.1030E+00	0.2133E+01	0.7952E+00	-.1396E+01	0.9885E+00	0.1929E+02
19	4.5000	0.6503E+00	0.2201E+01	-.2522E+00	-.8012E+00	0.2745E+01	-.5238E+01
20	4.7500	0.1181E+01	0.1991E+01	-.1429E+01	-.3879E+00	0.1236E+00	-.1573E+02
21	5.0000	0.1617E+01	0.1437E+01	-.3003E+01	-.6516E+00	-.1444E+01	0.3194E+01
22	5.2500	0.1874E+01	0.5849E+00	-.3813E+01	-.6909E+00	0.2016E+01	0.2449E+02
23	5.5000	0.1912E+01	-.2353E+00	-.2748E+01	0.4886E+00	0.7060E+01	0.1586E+02
24	5.7500	0.1792E+01	-.6351E+00	-.4500E+00	0.2425E+01	0.7132E+01	-.1527E+02
25	6.0000	0.1636E+01	-.5461E+00	0.1163E+01	0.3569E+01	0.1377E+01	-.3077E+02

Equation:  $5 \ddot{x} + 4 \dot{x} + 3x = 6 \sin t$

11.41 or  $\ddot{x} = -0.8 \dot{x} - 0.6x + 1.2 \sin t$

differential equations to be solved:

$$\frac{d\vec{Y}}{dt} = \begin{Bmatrix} y_2 \\ -0.8y_2 - 0.6y_1 + 1.2 \sin t \end{Bmatrix}$$


with  $\vec{Y} = \begin{Bmatrix} y_1 \\ y_2 \end{Bmatrix} = \begin{Bmatrix} x \\ \dot{x} \end{Bmatrix}$  and  $\vec{Y}(0) = \vec{0}$

```

% Ex11_41.m
% This program will use the function dfunc11_41.m, they should
% be in the same folder
tspan = [0: 0.1: 20];
x0 = [0.0; 0.0];
[t,x] = ode23('dfunc11_41', tspan, x0);
disp('      t      x(t)      xd(t)');
disp([t x]);
plot(t,x(:,1));
xlabel('t');
gtext('x(t)');

% dfunc11_41.m
function f = dfunc11_41(t,x)
f = zeros(2,1);
f(1) = x(2);
f(2) = - 0.8 * x(2) - 0.6 * x(1) + 1.2*sin(t);

```



Results of Ex11\_41

\*\*\*\*\*

>> Ex11\_41

t	x(t)	xd(t)
0	0	0
0.1000	0.0002	0.0058
0.2000	0.0015	0.0226
0.3000	0.0051	0.0493
0.4000	0.0117	0.0847
0.5000	0.0222	0.1274
0.6000	0.0374	0.1762
0.7000	0.0576	0.2298
0.8000	0.0835	0.2867
0.9000	0.1151	0.3456
1.0000	0.1526	0.4051

19.1000	-1.3092	-0.2840
19.2000	-1.3310	-0.1518
19.3000	-1.3395	-0.0181
19.4000	-1.3346	0.1158
19.5000	-1.3163	0.2486
19.6000	-1.2849	0.3789
19.7000	-1.2406	0.5054
19.8000	-1.1839	0.6268
19.9000	-1.1154	0.7420
20.0000	-1.0358	0.8498

11.42 Equations:  $2 \ddot{x}_1 + 10 x_1 - 5 x_2 = F_1(t)$   
 $4 \ddot{x}_2 - 5 x_1 + 15 x_2 = 0$


or  $\ddot{x}_1 = -5 x_1 + 2.5 x_2 + 0.5 F_1(t)$   
 $\ddot{x}_2 = 1.25 x_1 - 3.75 x_2$

Equations in vector form:

$$\frac{d\vec{Y}}{dt} = \begin{Bmatrix} y_2 \\ -5 y_1 + 2.5 y_3 + 0.5 F_1(t) \\ y_4 \\ 1.25 y_1 - 3.75 y_3 \end{Bmatrix} \equiv \vec{f}(t)$$

with

$$\vec{Y} = \begin{Bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{Bmatrix} = \begin{Bmatrix} x_1 \\ \dot{x}_1 \\ x_2 \\ \dot{x}_2 \end{Bmatrix}, \quad \vec{Y}(0) = \vec{0} \text{ and}$$




```
% Ex11_42.m
% This program will use the function dfunc11_42.m, they should
% be in the same folder
tspan = [0: 0.1: 20];
x0 = [0.0; 0.0; 0.0; 0.0];
[t,x] = ode23('dfunc11_42', tspan, x0);
disp('      t      x1(t)      xd1(t)      x2(t)      xd2(t)');
disp([t x]);
plot(t,x(:,1));
xlabel('t');
ylabel('Solid line: x_1(t) Dotted line: x_2(t)');
hold on;
plot(t,x(:,3), ':');
```

```

% dfunc11_42.m
function f = dfunc11_42(t,x)
F = 5*( stepfun(t,0) - stepfun(t,2.0) );
f = zeros(4,1);
f(1) = x(2);
f(2) = 2.5 * x(3) - 5 * x(1) + 0.5*F;
f(3) = x(4);
f(4) = 1.25 * x(1) - 3.75 * x(3);

```



Results of Ex11\_42

\*\*\*\*\*

>> Ex11\_42

t	x1(t)	xd1(t)	x2(t)	xd2(t)
0	0	0	0	0
0.1000	0.0124	0.2479	0.0000	0.0005
0.2000	0.0492	0.4835	0.0002	0.0041
0.3000	0.1084	0.6952	0.0010	0.0135
0.4000	0.1871	0.8725	0.0032	0.0311
0.5000	0.2814	1.0072	0.0076	0.0583
0.6000	0.3869	1.0932	0.0152	0.0958
0.7000	0.4984	1.1272	0.0271	0.1434
0.8000	0.6106	1.1086	0.0442	0.1995
0.9000	0.7184	1.0397	0.0672	0.2620
1.0000	0.8170	0.9256	0.0967	0.3277
19.3000	-0.1051	-0.9836	-0.5591	-0.7481
19.4000	-0.2073	-1.0540	-0.6238	-0.5451
19.5000	-0.3146	-1.0855	-0.6678	-0.3349
19.6000	-0.4229	-1.0711	-0.6907	-0.1256
19.7000	-0.5273	-1.0066	-0.6931	0.0750
19.8000	-0.6225	-0.8902	-0.6762	0.2603
19.9000	-0.7036	-0.7229	-0.6417	0.4248
20.0000	-0.7655	-0.5090	-0.5920	0.5644

11.43 Equations to be solved: (see Problems 11.18 and 11.29)

$$\frac{d\vec{Y}}{dt} = \begin{Bmatrix} y_2 \\ -2y_2 + 2y_4 - 6y_1 + 2y_3 \\ y_4 \\ y_2 - y_4 + y_1 - 4y_3 + 5 \end{Bmatrix}$$

with

$$\vec{Y} = \begin{Bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{Bmatrix} = \begin{Bmatrix} x_1 \\ \dot{x}_1 \\ x_2 \\ \dot{x}_2 \end{Bmatrix} \quad \text{and} \quad \vec{Y}(0) = \vec{0}$$

Results of Ex11\_43

\*\*\*\*\*

I	TIME(I)	X(1)	X(2)	X(3)	X(4)
1	0.15708000	0.00595234	0.11190540	0.07562662	0.69862971
2	0.31416000	0.04296424	0.38514540	0.25870383	1.23457026
3	0.47124000	0.12871802	0.72177873	0.51077890	1.61270950
4	0.62832000	0.26619304	1.03218582	0.79478663	1.85249641
5	0.78540000	0.44606930	1.24885462	1.07929835	1.98056156
6	0.94248000	0.65071547	1.33306426	1.34088682	2.02464088
7	1.09956000	0.85871211	1.27520063	1.56476458	2.00934956
8	1.25664000	1.04904086	1.09026067	1.74407153	1.95390776
9	1.41372000	1.20434253	0.81046444	1.87829054	1.87158084
10	1.57080000	1.31294336	0.47685982	1.97127049	1.77038826
11	1.72788000	1.36961694	0.13148283	2.02926358	1.65455622
12	1.88496000	1.37525431	-0.18884744	2.05926699	1.52621604
13	2.04204000	1.33574082	-0.45654785	2.06782932	1.38695120
14	2.19912000	1.26038610	-0.65503054	2.06036319	1.23893451
15	2.35620000	1.16023580	-0.77849419	2.04091265	1.08554048
16	2.51328000	1.04653025	-0.83026508	2.01226511	0.93144276
17	2.67036000	0.92948759	-0.82029318	1.97627325	0.78229624
18	2.82744000	0.81749686	-0.76237983	1.93425605	0.64415368
19	2.98452000	0.71672563	-0.67160439	1.88737286	0.52277900
20	3.14160000	0.63108587	-0.56226450	1.83689975	0.42300142
21	3.29868000	0.56246606	-0.44648447	1.78437494	0.34821759
22	3.45576000	0.51112440	-0.33350502	1.73161322	0.30010274
23	3.61284000	0.47614448	-0.22956211	1.68061392	0.27854652
24	3.76992000	0.45587455	-0.13819995	1.63340072	0.28179218
25	3.92700000	0.44829676	-0.06084131	1.59183581	0.30673179
26	4.08408000	0.45130011	0.00254980	1.55744649	0.34929734
27	4.24116000	0.46285317	0.05284442	1.53129308	0.40488652
28	4.39824000	0.48108992	0.09124953	1.51389460	0.46876999
29	4.55532000	0.50433136	0.11897352	1.50521674	0.53644001
30	4.71240000	0.53106884	0.13705660	1.50471659	0.60387671
31	4.86948000	0.55993248	0.14633820	1.51143126	0.66772319
32	5.02656000	0.58966273	0.14751780	1.52409402	0.72537321
33	5.18364000	0.61909585	0.14126165	1.54126111	0.77498438
34	5.34072000	0.64716684	0.12831325	1.56143405	0.81543446
35	5.49780000	0.67292815	0.10957633	1.58316615	0.84623969
36	5.65488000	0.69557810	0.08615286	1.60514603	0.86745270
37	5.81196000	0.71449180	0.05933176	1.62625520	0.87955395
38	5.96904000	0.72924718	0.03053440	1.64560020	0.88334709
39	6.12612000	0.73964026	0.00123034	1.66252252	0.87986403
40	6.28320000	0.74568650	-0.02716047	1.67659091	0.87028266

11.44 Equations of motion:

$$\begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} \ddot{\vec{x}} + \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \dot{\vec{x}} + \begin{bmatrix} 6 & -2 \\ -2 & 8 \end{bmatrix} \vec{x} = \vec{f} = \begin{Bmatrix} 10 \sin 5t \\ 0 \end{Bmatrix}$$

with  $\vec{x}(0) = \vec{0}$  and  $\dot{\vec{x}}(0) = \vec{0}$

Program 15.cpp is used for solution.

Results of Ex11\_44

\*\*\*\*\*

Please input N and NSTEP:

2 24

Please input M matrix row by row

1 0

0 2

Please input K matrix row by row

6 -2

-2 8

Please input C matrix row by row

0 0

0 0

SOLUTION BY CENTRAL DIFFERENCE METHOD

GIVEN DATA:

N = 2 NSTEP = 24 DELT = 0.24216267

SOLUTION:

STEP	TIME	X(I,1)	XD(I,1)	XDD(I,1)	X(I,2)	XD(I,2)	XDD(I,2)
1	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	5.0000
2	0.2422	0.0000	0.0000	0.0000	0.1466	0.0000	5.0000
3	0.4843	0.0172	0.0355	0.2932	0.5520	1.1398	4.4136
4	0.7265	0.0931	0.1922	1.0009	1.1222	2.0143	2.8090
5	0.9687	0.2678	0.5175	1.6859	1.7278	2.4276	0.6043
6	1.2108	0.5510	0.9455	1.8485	2.2370	2.3018	-1.6433
7	1.4530	0.9027	1.3107	1.1680	2.5470	1.6915	-3.3971
8	1.6951	1.2354	1.4132	-0.3219	2.6057	0.7613	-4.2855
9	1.9373	1.4391	1.1077	-2.2012	2.4189	-0.2646	-4.1875
10	2.1795	1.4202	0.3814	-3.7971	2.0422	-1.1635	-3.2363
11	2.4216	1.1410	-0.6155	-4.4366	1.5630	-1.7671	-1.7487
12	2.6638	0.6437	-1.6032	-3.7201	1.0773	-1.9923	-0.1110
13	2.9060	0.0463	-2.2604	-1.7077	0.6698	-1.8441	1.3345
14	3.1481	-0.4889	-2.3385	1.0621	0.4012	-1.3960	2.3669
15	3.3903	-0.8050	-1.7576	3.7357	0.3030	-0.7575	2.9063
16	3.6324	-0.8023	-0.6471	5.4359	0.3797	-0.0444	2.9831
17	3.8746	-0.4728	0.6859	5.5732	0.6135	0.6412	2.6789
18	4.1168	0.0950	1.8528	4.0638	0.9689	1.2166	2.0731
19	4.3589	0.7431	2.5104	1.3676	1.3958	1.6152	1.2194
20	4.6011	1.2933	2.4742	-1.6667	1.8321	1.7822	0.1598
21	4.8433	1.6034	1.7764	-4.0959	2.2077	1.6763	-1.0350
22	5.0854	1.6083	0.6502	-5.2053	2.4526	1.2813	-2.2272
23	5.3276	1.3349	-0.5545	-4.7444	2.5098	0.6238	-3.2023
24	5.5697	0.8862	-1.4909	-2.9897	2.3498	-0.2124	-3.7043
25	5.8119	0.4013	-1.9277	-0.6176	1.9837	-1.0863	-3.5128

11.45

Equations of motion: (See Problems 11.20 and 11.31)

$$\begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix} \ddot{\vec{x}} + \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \dot{\vec{x}} + \begin{bmatrix} 6 & -2 \\ -2 & 4 \end{bmatrix} \vec{x} = \vec{F}(t) = \begin{Bmatrix} 5 \\ 20 \sin 5t \end{Bmatrix}$$

$$\text{with } \vec{x}(0) = \vec{0} \text{ and } \dot{\vec{x}}(0) = \vec{0}$$

Results of Ex11\_45

\*\*\*\*\*

Please input N and NSTEP:

2 50

Please input M matrix row by row

2 0

0 1

Please input K matrix row by row

6 -2

-2 4

Please input C matrix row by row

0 0

0 0

SOLUTION BY HOUBOLT METHOD

GIVEN DATA:

N = 2 NSTEP = 50 DELT = 0.25000000

SOLUTION:

STEP	TIME	X(I,1)	XD(I,1)	XDD(I,1)	X(I,2)	XD(I,2)	XDD(I,2)
1	0.0000	0.0000	0.0000	2.5000	0.0000	0.0000	0.0000
2	0.2500	0.0781	0.0000	2.5000	0.0000	0.0000	0.0000
3	0.5000	0.2979	0.5957	2.2656	1.1960	2.3920	19.1359
4	0.7500	0.7015	2.0388	3.6195	3.2240	9.2904	7.4868
5	1.0000	1.3214	2.9557	3.9797	5.4440	8.4114	-7.1638
6	1.2500	2.1367	3.6237	2.7889	6.6991	1.5474	-33.9541
7	1.5000	3.0390	3.6390	-0.3395	6.2776	-5.9881	-38.2107
8	1.7500	3.8478	2.8069	-4.3877	4.6555	-8.2542	-11.5902
9	2.0000	4.3696	1.2560	-7.6821	2.9268	-5.6702	15.7921
10	2.2500	4.4586	-0.7048	-9.2635	1.6122	-3.7352	14.9627
11	2.5000	4.0419	-2.7752	-9.2545	0.3710	-5.2727	-4.2808
12	2.7500	3.1231	-4.6742	-7.9744	-1.1051	-6.7851	-8.6896
13	3.0000	1.7906	-6.0399	-5.2080	-2.3361	-3.7943	11.5993
14	3.2500	0.2287	-6.4607	-0.7209	-2.5348	2.3200	29.1162
15	3.5000	-1.3005	-5.7014	4.7195	-1.6819	5.5404	17.1324
16	3.7500	-2.5205	-3.8938	9.3655	-0.6961	2.9838	-12.5743
17	4.0000	-3.2188	-1.4660	11.7521	-0.4043	-1.3232	-24.3329
18	4.2500	-3.2899	1.1109	11.7232	-0.6464	-1.8229	-5.9818
19	4.5000	-2.7339	3.4779	10.0298	-0.6718	1.3328	15.4785
20	4.7500	-1.6400	5.3326	7.1804	-0.2395	2.9664	11.1806
21	5.0000	-0.1792	6.3485	3.1320	0.0943	0.3961	-10.4792
22	5.2500	1.4071	6.2745	-1.8521	-0.1309	-2.6326	-16.3098
23	5.5000	2.8519	5.1403	-6.5357	-0.4800	-1.0645	4.9768
24	5.7500	3.9330	3.3011	-9.3734	-0.0742	4.3048	26.1411
:							
46	11.2500	3.7024	0.3659	-4.9785	3.6288	-2.1394	-13.8351
47	11.5000	3.6344	-0.9623	-5.8596	2.5437	-6.8485	-22.9011
48	11.7500	3.2197	-2.4109	-6.2469	0.9121	-7.1488	-3.0939
49	12.0000	2.4690	-3.6600	-5.2000	-0.2931	-2.6700	22.3940
50	12.2500	1.4714	-4.3654	-2.5259	-0.6117	1.1118	21.5438
51	12.5000	0.3752	-4.3846	0.7930	-0.5814	0.1019	-3.0201

11.46

The program and output are given below.

```

DIMENSION X(4), XT(4), XTN(4), FT(4), A(4, 4), F(2), M(2, 2), K(2, 2),
2 C(2, 2), MI(2, 2), MIK(2, 2), MIC(2, 2), MIF(2), ZK1(4), ZK2(4),
3 ZK3(4), ZK4(4), XD(2), XDD(2), XX(2), X1(2), X2(2)
REAL M, K, MI, MIK, MIC, MIF
DATA N, NSTEP, T, DELT/2, 25, 0, 0, 0, 24216267/
DATA X/0., 0., 0., 0. /
DATA M/1., 0., 0., 2. /
DATA C/1. 0, -1. 0, -1. 0, 1. 0/
DATA K/6., -2., -2., 8. /
N2=N*2
DO 11 I=1, NSTEP
CALL RKM4 (N, N2, T, DELT, X, XT, XTN, FT, A, F, M, K, C, MI, MIK, MIC,
2 MIF, ZK1, ZK2, ZK3, ZK4)
DO 13 II=1, N
XX(II)=X(II)
13 XD(II)=X(II+N)
CALL XMULT (MIC, XD, X1, N)
CALL XMULT (MIK, XX, X2, N)
DO 14 II=1, N
14 XDD(II)=MIF(II)-X1(II)-X2(II)
PRINT 12, I, T, ((XX(J), XD(J), XDD(J)), J=1, N)
12 FORMAT (2X, I2, 2X, F6. 3, 1X, 3E10. 2, 1X, 3E10. 2)
11 CONTINUE
STOP
END

```

```

C
SUBROUTINE RKM4 (N, N2, T, DELT, X, XT, XTN, FT, A, F, M, K, C, MI,
2 MIK, MIC, MIF, ZK1, ZK2, ZK3, ZK4)
REAL M, K, MI, MIK, MIC, MIF
DIMENSION X(N2), XT(N2), XTN(N2), FT(N2), A(N2, N2), F(N), M(N, N),
2 K(N, N), C(N, N), MI(N, N), MIK(N, N), MIC(N, N), MIF(N),
3 ZK1(N2), ZK2(N2), ZK3(N2), ZK4(N2)
DO 11 I=1, N2
11 XT(I)=X(I)
TIME=T
CALL FUN (N, N2, XT, FT, TIME, A, F, M, K, C, MI, MIK, MIC, MIF)
DO 12 I=1, N2
12 ZK1(I)=DELT*FT(I)

```

```

DO 13 I=1,N2
13 XT(I)=X(I)+0.5*ZK1(I)
   TIME=T+0.5*DELT
   CALL FUN (N,N2,XT,FT,TIME,A,F,M,K,C,MI,MIK,MIC,MIF)
   DO 14 I=1,N2
14   ZK2(I)=DELT*FT(I)
      DO 15 I=1,N2
15     XT(I)=X(I)+0.5*ZK2(I)
        CALL FUN (N,N2,XT,FT,TIME,A,F,M,K,C,MI,MIK,MIC,MIF)
        DO 16 I=1,N2
16       ZK3(I)=DELT*FT(I)
          DO 17 I=1,N2
17         XT(I)=X(I)+ZK3(I)
           TIME=T+DELT
           CALL FUN (N,N2,XT,FT,TIME,A,F,M,K,C,MI,MIK,MIC,MIF)
           DO 18 I=1,N2
18           ZK4(I)=DELT*FT(I)
             DO 19 I=1,N2
19             XTN(I)=X(I)+(ZK1(I)+2.0*ZK2(I)+2.0*ZK3(I)+ZK4(I))/6.0
               DO 21 I=1,N2
21               X(I)=XTN(I)
                 T=TIME
                 RETURN
                 END
                 SUBROUTINE FUN (N,N2,X,FT,T,A,F,M,K,C,MI,MIK,MIC,MIF)
                   REAL M,K,MI,MIK,MIC,MIF
                   DIMENSION X(N2),A(N2,N2),FT(N2),M(N,N),K(N,N),C(N,N),MI(N,N),
2 MIK(N,N),MIC(N,N),MIF(N),F(N)
C INVERSE OF M-MATRIX
   DET=M(1,1)*M(2,2)-M(1,2)*M(2,1)
   MI(1,1)=M(2,2)/DET
   MI(1,2)=-M(1,2)/DET
   MI(2,1)=-M(2,1)/DET
   MI(2,2)=M(1,1)/DET
C END OF INVERSE OF M-MATRIX
   CALL MATMUL (MIK,MI,K,N,N,N)
   CALL MATMUL (MIC,MI,C,N,N,N)
C DEFINITION OF F-VECTOR
   CALL FUN2 (X,T,N,N2,F)
C END OF F-VECTOR
   CALL XMULT (MI,F,MIF,N)
   DO 11 I=1,N
   DO 11 J=1,N
   A(I,J)=0.0
   A(I,J+N)=0.0
   A(I+N,J)=-MIK(I,J)
11  A(I,J+N)=-MIC(I,J)
   DO 12 I=1,N
12  A(I,I+N)=1.0
   CALL XMULT (A,X,FT,N2)
   N1=N+1
   DO 13 I=N1,N2
13  FT(I)=FT(I)+MIF(I-N)
   RETURN
   END

```

```

C
SUBROUTINE FUN2 (X, T, N, N2, F)
DIMENSION X(N2), F(N)
F(1)=0.0
F(2)=10.0
RETURN
END

```

```

C
SUBROUTINE XMULT (A, B, BB, N)
DIMENSION A(N, N), B(N), BB(N)
DO 10 I=1, N
BB(I)=0.0
DO 10 J=1, N
10 BB(I)=BB(I)+A(I, J)*B(J)
RETURN
END

```

```

C
SUBROUTINE MATMUL (A, B, C, L, M, N)
DIMENSION A(L, N), B(L, M), C(M, N)
DO 10 I=1, L
DO 10 J=1, N
A(I, J)=0.0
DO 10 K=1, M
10 A(I, J)=A(I, J)+B(I, K)*C(K, J)
RETURN
END

```

$i$	$t_i$	$x_{1i}$	$\dot{x}_{1i}$	$\ddot{x}_{1i}$	$x_{2i}$	$\dot{x}_{2i}$	$\ddot{x}_{2i}$
1	0.242	0.14E+00	-0.47E-01	0.66E+00	0.14E+00	0.12E+01	0.40E+01
2	0.484	0.51E+00	-0.35E+00	0.50E+00	0.53E+00	0.22E+01	0.21E+01
3	0.726	0.95E+00	-0.10E+01	0.19E+00	0.11E+01	0.28E+01	-0.17E+00
4	0.969	0.13E+01	-0.20E+01	0.43E+00	0.16E+01	0.30E+01	-0.75E+01
5	1.211	0.14E+01	-0.31E+01	0.17E+01	0.20E+01	0.28E+01	-0.45E+01
6	1.453	0.11E+01	-0.40E+01	0.39E+01	0.22E+01	0.23E+01	-0.56E+01
7	1.695	0.59E+00	-0.42E+01	0.66E+01	0.21E+01	0.16E+01	-0.58E+01
8	1.937	-0.66E-01	-0.36E+01	0.88E+01	0.20E+01	0.92E+00	-0.51E+01
9	2.179	-0.64E+00	-0.22E+01	0.97E+01	0.17E+01	0.25E+00	-0.38E+01
10	2.422	-0.94E+00	-0.18E+00	0.86E+01	0.16E+01	-0.33E+00	-0.22E+01
11	2.664	-0.87E+00	0.19E+01	0.59E+01	0.15E+01	-0.86E+00	-0.66E+00
12	2.906	-0.44E+00	0.37E+01	0.88E+00	0.16E+01	-0.13E+01	0.57E+00
13	3.148	0.21E+00	0.47E+01	-0.42E+01	0.18E+01	-0.18E+01	0.14E+01
14	3.390	0.88E+00	0.48E+01	-0.86E+01	0.19E+01	-0.22E+01	0.19E+01
15	3.632	0.14E+01	0.40E+01	-0.11E+02	0.18E+01	-0.25E+01	0.24E+01
16	3.875	0.16E+01	0.26E+01	-0.11E+02	0.16E+01	-0.26E+01	0.28E+01
17	4.117	0.14E+01	0.11E+01	-0.95E+01	0.12E+01	-0.24E+01	0.33E+01
18	4.359	0.99E+00	-0.17E+00	-0.61E+01	0.74E+00	-0.18E+01	0.39E+01
19	4.601	0.50E+00	-0.10E+01	-0.23E+01	0.32E+00	-0.96E+00	0.42E+01
20	4.843	0.11E+00	-0.13E+01	0.99E+00	0.79E-01	0.15E+00	0.41E+01
21	5.085	-0.39E-01	-0.13E+01	0.30E+01	0.93E-01	0.13E+01	0.33E+01
22	5.328	0.97E-01	-0.12E+01	0.37E+01	0.38E+00	0.23E+01	0.18E+01
23	5.570	0.45E+00	-0.13E+01	0.34E+01	0.88E+00	0.30E+01	-0.21E+00
24	5.812	0.85E+00	-0.17E+01	0.27E+01	0.15E+01	0.32E+01	-0.24E+01
25	6.054	0.11E+01	-0.23E+01	0.25E+01	0.20E+01	0.30E+01	-0.44E+01

11.47

```

C =====
C
C PROGRAM 15.F
C MAIN PROGRAM WHICH CALLS CDIFF
C
C =====
C FOLLOWING 10 LINES CONTAIN PROBLEM-DEPENDENT DATA
  REAL M(2,2),K(2,2),MC(2,2),MK(2,2),MCI(2,2),MMC(2,2),MI(2,2)
  DIMENSION C(2,2),XI(2),XDI(2),XDDI(2),XM1(2),F(2),R(2),RR(2),
  2  XMK(2),XMI(2),XM2(2),XP1(2),ZA(2),ZB(2),ZC(2),LA(2),LB(2,2),
  3  S(2),X(25,2),XD(25,2),XDD(25,2)
  DATA N,NSTEP,NSTEP1,DELT/2,24,25,0.24216267/
  DATA XI/0.0,0.0/
  DATA XDI/0.0,0.0/
  DATA M/1.0,0.0,0.0,2.0/
  DATA C/0.0,0.0,0.0,0.0/
  DATA K/6.0,-2.0,-2.0,8.0/
C END OF PROBLEM-DEPENDENT DATA
  CALL CDIFF (M,C,K,XI,XDI,XDDI,N,NSTEP,DELT,F,R,RR,XM1,XM2,XP1,
  2  MC,MK,MCI,XMK,MMC,XMI,ZA,ZB,ZC,LA,LB,S,X,XD,XDD,NSTEP1,MI)
  PRINT 10
  10  FORMAT (//,38H SOLUTION BY CENTRAL DIFFERENCE METHOD,/)
  PRINT 20, N,NSTEP,DELT
  20  FORMAT (12H GIVEN DATA: ,/,3H N=,I5,4X,7H NSTEP=,I5,4X,6H DELT=,
  2  E15.8,/)
  PRINT 30
  30  FORMAT (10H SOLUTION: ,/,5H STEP,3X,5H TIME,3X,7H X(I,1),3X,
  2  8H XD(I,1),2X,9H XDD(I,1),4X,7H X(I,2),3X,8H XD(I,2),2X,
  3  9H XDD(I,2),/)
  DO 40, I=1,NSTEP1
  TIME=REAL(I-1)*DELT
  40  PRINT 50, I,TIME,X(I,1),XD(I,1),XDD(I,1),X(I,2),XD(I,2),XDD(I,2)
  50  FORMAT (1X,I4,F8.4,6(1X,E10.4))
  STOP
  END
C =====
C
C SUBROUTINE CDIFF
C
C =====
  SUBROUTINE CDIFF (M,C,K,XI,XDI,XDDI,N,NSTEP,DELT,F,R,RR,XM1,
  2  XM2,XP1,MC,MK,MCI,XMK,MMC,XMI,ZA,ZB,ZC,LA,LB,S,X,XD,XDD,NSTEP1
  3  ,MI)
  REAL M(N,N),K(N,N),MC(N,N),MK(N,N),MCI(N,N),MMC(N,N),MI(N,N)
  DIMENSION C(N,N),XI(N),XDI(N),XDDI(N),F(N),R(N),RR(N),XM1(N),
  2  XM2(N),XP1(N),XMK(N),XMI(N),ZA(N),ZB(N),ZC(N),LA(N),LB(N,N),S(N)
  3  ,X(NSTEP1,N),XD(NSTEP1,N),XDD(NSTEP1,N)
  DO 5 I=1,N
  DO 5 J=1,N
  5  MI(I,J)=M(I,J)
  CALL SIMUL (MI,ZA,N,0,LA,LB,S)
  CALL EXTFUN (F,0.0,N)
  DO 20 I=1,N
  R(I)=F(I)
  DO 10 J=1,N
  10  R(I)=R(I)-C(I,J)*XDI(J)-K(I,J)*XI(J)
  20  CONTINUE
  CALL XMULT (MI,R,XDDI,N)
  DO 25 J=1,N
  X(1,J)=XI(J)
  XD(1,J)=XDI(J)
  25  XDD(1,J)=XDDI(J)
  DO 30 I=1,N
  30  XM1(I)=XI(I)-DELT*XDI(I)+(DELT**2)*XDDI(I)/2.0
  DO 40 I=1,N
  DO 40 J=1,N
  MC(I,J)=(M(I,J)/(DELT**2))+(C(I,J)/(2.0*DELT))
  MK(I,J)=K(I,J)-2.0*M(I,J)/(DELT**2)
  40  MMC(I,J)=(M(I,J)/(DELT**2))-(C(I,J)/(2.0*DELT))
  DO 45 I=1,N
  DO 45 J=1,N
  45  MCI(I,J)=MC(I,J)

```

```

CALL SIMUL (MCI,ZA,N,0,LA,LB,S)
TIME=DELT
DO 90 I=1,NSTEP
CALL XMULT (MK,XI,XMK,N)
CALL XMULT (MMC,XM1,XMI,N)
TIME=TIME+DELT
CALL EXTFUN (F,TIME,N)
DO 50 J=1,N
50 RR(J)=F(J)-XMK(J)-XMI(J)
CALL XMULT (MCI,RR,XP1,N)
DO 60 J=1,N
XDI(J)=(XP1(J)-XM1(J))/(2.0*DELT)
60 XDDI(J)=(XP1(J)-2.0*XI(J)+XM1(J))/(DELT**2)
DO 70 J=1,N
XM1(J)=XI(J)
70 XI(J)=XP1(J)
DO 80 J=1,N
X(I+1,J)=XP1(J)
XD(I+1,J)=XDI(J)
80 XDD(I+1,J)=XDDI(J)
90 CONTINUE
RETURN
END
C =====
C
C SUBROUTINE EXTFUN
C THIS SUBROUTINE IS PROBLEM-DEPENDENT
C
C =====
SUBROUTINE EXTFUN (F,TIME,N)
DIMENSION F(N)
F(1) = 10.0*SIN(5.0*TIME)
F(2) = 0.0
RETURN
END
C =====
C
C SUBROUTINE XMULT
C
C =====
SUBROUTINE XMULT (A,B,BB,N)
DIMENSION A(N,N),B(N),BB(N)
DO 10 I=1,N
BB(I)=0.0
DO 10 J=1,N
10 BB(I)=BB(I)+A(I,J)*B(J)
RETURN
END
C =====
C
C SUBROUTINE SIMUL
C
C =====
SUBROUTINE SIMUL (A,B,N,IND,LA,LB,S)
DIMENSION A(N,N),B(N),LA(N),LB(N,2),S(N)
DO 100 I=1,N
100 LA(I)=0
DO 250 K=1,N
Z=0.0
DO 150 I=1,N
IF (LA(I) .EQ. 1) GO TO 150
DO 140 J=1,N
IF (LA(J)-1) 130,140,300
130 IF (ABS(Z) .GE. ABS(A(I,J))) GO TO 140
IA=I
IB=J
Z=A(I,J)
140 CONTINUE
150 CONTINUE

```

```

      LA(IB)=LA(IB)+1
      IF (IA .EQ. IB) GO TO 190
      DO 160 I=1,N
      Z=A(IA,I)
      A(IA,I)=A(IB,I)
160   A(IB,I)=Z
      IF (IND .EQ. 0) GO TO 190
      Z=B(IA)
      B(IA)=B(IB)
      B(IB)=Z
190   LB(K,1)=IA
      LB(K,2)=IB
      S(K)=A(IB,IB)
      A(IB,IB)=1.0
      DO 200 I=1,N
200   A(IB,I)=A(IB,I)/S(K)
      IF (IND .EQ. 0) GO TO 220
      B(IB)=B(IB)/S(K)
220   DO 250 I=1,N
      IF(I .EQ. IB) GO TO 250
      Z=A(I,IB)
      A(I,IB)=0.0
      DO 230 J=1,N
230   A(I,J)=A(I,J)-A(IB,J)*Z
      IF (IND .EQ. 0) GO TO 250
      B(I)=B(I)-B(IB)*Z
250   CONTINUE
      DO 270 I=1,N
      J=N-I+1
      IF (LB(J,1) .EQ. LB(J,2)) GO TO 270
      IA=LB(J,1)
      IB=LB(J,2)
      DO 260 K=1,N
      Z=A(K,IA)
      A(K,IA)=A(K,IB)
      A(K,IB)=Z
260   CONTINUE
270   CONTINUE
300   RETURN
      END

```

## SOLUTION BY CENTRAL DIFFERENCE METHOD

## GIVEN DATA:

N= 2      NSTEP= 24      DELT= 0.24216267E+00

## SOLUTION:

STEP	TIME	X(I,1)	XD(I,1)	XDD(I,1)	X(I,2)	XD(I,2)	XDD(I,2)
1	0.0000	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00
2	0.2422	0.3867E+00	0.7984E+00	0.6594E+01	0.0000E+00	0.0000E+00	0.0000E+00
3	0.4843	0.3609E+00	0.7451E+00	-.7034E+01	0.2268E-01	0.4682E-01	0.3867E+00
4	0.7265	-.3707E+00	-.1564E+01	-.1203E+02	0.6119E-01	0.1263E+00	0.2702E+00
5	0.9687	-.1098E+01	-.3012E+01	0.7527E-01	0.6362E-01	0.8454E-01	-.6155E+00
6	1.2108	-.9436E+00	-.1183E+01	0.1503E+02	-.1326E-01	-.1537E+00	-.1352E+01
7	1.4530	0.1774E-01	0.2303E+01	0.1376E+02	-.1424E+00	-.4253E+00	-.8906E+00
8	1.6951	0.8044E+00	0.3609E+01	-.2979E+01	-.2370E+00	-.4620E+00	0.5872E+00
9	1.9373	0.6966E+00	0.1402E+01	-.1525E+02	-.2289E+00	-.1787E+00	0.1753E+01
10	2.1795	0.5749E-01	-.1542E+01	-.9061E+01	-.1263E+00	0.2287E+00	0.1612E+01
11	2.4216	-.2159E+00	-.1884E+01	0.6238E+01	0.9378E-02	0.4920E+00	0.5626E+00
12	2.6638	0.1297E+00	0.1490E+00	0.1055E+02	0.1302E+00	0.5295E+00	-.2534E+00
13	2.9060	0.4257E+00	0.1325E+01	-.8437E+00	0.2280E+00	0.4514E+00	-.3910E+00
14	3.1481	0.4341E-01	-.1781E+00	-.1157E+02	0.2974E+00	0.3452E+00	-.4864E+00
15	3.3903	-.6914E+00	-.2306E+01	-.6011E+01	0.2995E+00	0.1475E+00	-.1146E+01
16	3.6324	-.8546E+00	-.1854E+01	0.9746E+01	0.1908E+00	-.2200E+00	-.1889E+01
17	3.8746	-.1162E+00	0.1188E+01	0.1538E+02	-.1272E-01	-.6446E+00	-.1618E+01
18	4.1168	0.7762E+00	0.3367E+01	0.2624E+01	-.2201E+00	-.8484E+00	-.6526E-01
19	4.3589	0.8716E+00	0.2040E+01	-.1359E+02	-.3303E+00	-.6557E+00	0.1657E+01
20	4.6011	0.1564E+00	-.1280E+01	-.1383E+02	-.3119E+00	-.1897E+00	0.2193E+01

```

21 4.8433 -.4804E+00 -.2792E+01 0.1339E+01 -.2112E+00 0.2459E+00 0.1404E+01
22 5.0854 -.3877E+00 -.1123E+01 0.1244E+02 -.8914E-01 0.4600E+00 0.3645E+00
23 5.3276 0.7305E-01 0.1143E+01 0.6277E+01 0.3113E-01 0.5004E+00 -.3116E-01
24 5.5697 0.9719E-01 0.1001E+01 -.7446E+01 0.1484E+00 0.4904E+00 -.5145E-01
25 5.8119 -.4297E+00 -.1038E+01 -.9396E+01 0.2365E+00 0.4241E+00 -.4963E+00

```

11.48 The data to be used in the main program which calls HOBOLT, subroutine EXTFUN and the output are given below.

C FOLLOWING 10 LINES CONTAIN PROBLEM-DEPENDENT DATA

```

REAL M(2,2),K(2,2),MC(2,2),MK(2,2),MCI(2,2),M*MC(2,2)
DIMENSION C(2,2),XI(2),XDI(2),XDDI(2),XMI(2),F(2),R(2),RR(2),
2 XMK(2),XMI(2),XM2(2),XPI(2),ZA(2),ZB(2),ZC(2),LA(2),LB(2,2),
3 S(2),X(25,2),XD(25,2),XDD(25,2)
DATA N,NSTEP,NSTEP1,DELT/2,24,25,0.25/

```

```

DATA XI/0.0,0.0/
DATA XDI/0.0,0.0/
DATA M/2.0,0.0,0.0,1.0/
DATA C/0.0,0.0,0.0,0.0/
DATA K/6.0,-2.0,-2.0,4.0/

```

C END OF PROBLEM-DEPENDENT DATA

```

SUBROUTINE EXTFUN (F,TIME,N)
DIMENSION F(N)
F(1)=5.0
F(2)=20.0*SIN(5.0*TIME)
RETURN
END

```

SOLUTION BY HOBOLT METHOD

GIVEN DATA:

N = 2    NSTEP = 24    DELT = 0.25000000E+00

SOLUTION:

STEP	TIME	X(I,1)	XD(I,1)	XDD(I,1)	X(I,2)	XD(I,2)	XDD(I,2)
1	0.0000	0.0000E+00	0.0000E+00	0.2500E+01	0.0000E+00	0.0000E+00	0.0000E+00
2	0.2500	0.7813E-01	0.0000E+00	0.2500E+01	0.1186E+01	0.2372E+01	0.1898E+02
3	0.5000	0.3720E+00	0.7440E+00	0.3452E+01	0.2834E+01	0.5668E+01	0.7381E+01
4	0.7500	0.8908E+00	0.2538E+01	0.3748E+01	0.3920E+01	0.1861E+01	-.2533E+02
5	1.0000	0.1570E+01	0.2954E+01	0.1544E+01	0.3756E+01	-.4081E+01	-.3106E+02
6	1.2500	0.2281E+01	0.2728E+01	-.1597E+01	0.2744E+01	-.5200E+01	-.7080E+01
7	1.5000	0.2872E+01	0.1932E+01	-.4272E+01	0.1845E+01	-.2097E+01	0.1713E+02
8	1.7500	0.3225E+01	0.7715E+00	-.5759E+01	0.1416E+01	-.2943E+00	0.1328E+02
9	2.0000	0.3261E+01	-.5946E+00	-.6317E+01	0.9659E+00	-.2500E+01	-.8222E+01
10	2.2500	0.2943E+01	-.2030E+01	-.6253E+01	0.7463E-01	-.5007E+01	-.1377E+02
11	2.5000	0.2285E+01	-.3287E+01	-.5188E+01	-.8318E+00	-.3088E+01	0.6572E+01
12	2.7500	0.1378E+01	-.4014E+01	-.2594E+01	-.9609E+00	0.2095E+01	0.2512E+02
13	3.0000	0.3796E+00	-.3959E+01	0.1120E+01	-.2410E+00	0.4673E+01	0.1473E+02
14	3.2500	-.5214E+00	-.3160E+01	0.4552E+01	0.4875E+00	0.1810E+01	-.1331E+02
15	3.5000	-.1171E+01	-.1890E+01	0.6491E+01	0.4775E+00	-.2512E+01	-.2376E+02
16	3.7500	-.1480E+01	-.4332E+00	0.6889E+01	-.4997E-01	-.2951E+01	-.4747E+01
17	4.0000	-.1417E+01	0.1036E+01	0.6435E+01	-.3148E+00	0.5060E+00	0.1668E+02
18	4.2500	-.9987E+00	0.2360E+01	0.5417E+01	-.7897E-01	0.2262E+01	0.1182E+02
19	4.5000	-.2950E+00	0.3294E+01	0.3460E+01	0.7496E-01	-.3250E+00	-.1063E+02
20	4.7500	0.5658E+00	0.3586E+01	0.4555E+00	-.3473E+00	-.3500E+01	-.1713E+02

```

21 5.0000 0.1421E+01 0.3191E+01 -.2697E+01 -.9347E+00 -.2132E+01 0.3933E+01
22 5.2500 0.2126E+01 0.2332E+01 -.4688E+01 -.8089E+00 0.3101E+01 0.2547E+02
23 5.5000 0.2599E+01 0.1315E+01 -.5055E+01 0.2427E+00 0.6341E+01 0.1821E+02
24 5.7500 0.2816E+01 0.3215E+00 -.4484E+01 0.1464E+01 0.4216E+01 -.9383E+01
25 6.0000 0.2786E+01 -.6007E+00 -.3789E+01 0.2068E+01 0.1335E+00 -.2246E+02

```

11.49

```

=====
C MAIN PROGRAM WHICH CALLS WILSON
C
C
C =====

```

```

C FOLLOWING 10 LINES CONTAIN PROBLEM-DEPENDENT DATA

```

```

REAL M(2,2),MI(2,2),K(2,2)
DIMENSION C(2,2),XI(2),XDI(2),XDDI(2),X(25,2),XD(25,2),XDD(25,2),
2 XI(2),F(2),F1(2),F2(2),FT(2),R(2),LA(2),LB(2,2),S(2),ZA(2),
3 TK(2,2),XN1(2),XN2(2),XN3(2),XN4(2)
DATA N,NSTEP,NSTEP1,TH,DELTA/2,24,25,1.4,0.24216267/
DATA XI/0.0,0.0/
DATA XDI/0.0,0.0/
DATA M/1.0,0.0,0.0,2.0/
DATA C/0.0,0.0,0.0,0.0/
DATA K/6.0,-2.0,-2.0,8.0/

```

```

C END OF PROBLEM-DEPENDENT DATA

```

```

CALL WILSON (M,C,K,MI,XI,XDI,XDDI,X,XD,XDD,XT,F,F1,F2,FT,R,LA,
2 LB,S,ZA,TK,XN1,XN2,XN3,XN4,NSTEP1,DELTA,TH,N)
PRINT 10
10 FORMAT (//,26H SOLUTION BY WILSON METHOD,/)
PRINT 20, N,NSTEP,TH,DELTA
20 FORMAT (12H GIVEN DATA: ,/,3H N=,15,4X,7H NSTEP=,15,4X,4H TH=,
2 E15.8,4X,7H DELTA=,E15.8,/)
PRINT 30
30 FORMAT (10H SOLUTION: ,//,5H STEP,3X,5H TIME,3X,7H X(I,1),3X,
2 8H XD(I,1),2X,9H XDD(I,1),4X,7H X(I,2),3X,8H XD(I,2),2X,
3 9H XDD(I,2),/)
DO 40 I=1,NSTEP1
TIME=REAL(I-1)*DELTA
40 PRINT 50, I,TIME,X(I,1),XD(I,1),XDD(I,1),X(I,2),XD(I,2),XDD(I,2)
50 FORMAT (1X,I4,F8.4,6(1X,E10.4))
STOP
END

```

```

C =====

```

```

C SUBROUTINE WILSON
C
C
C =====

```

```

SUBROUTINE WILSON (M,C,K,MI,XI,XDI,XDDI,X,XD,XDD,XI,F,F1,F2,FT,
2 R,LA,LB,S,ZA,TK,XN1,XN2,XN3,XN4,NSTEP1,DELTA,TH,N)
REAL M(N,N),MI(N,N),K(N,N)
DIMENSION C(N,N),XI(N),XDI(N),XDDI(N),X(NSTEP1,N),XD(NSTEP1,N),
2 XDD(NSTEP1,N),XT(N),F(N),F1(N),F2(N),FT(N),R(N),LA(N),LB(N,2),
3 S(N),ZA(N),TK(N,N),XN1(N),XN2(N),XN3(N),XN4(N)
DO 5 I=1,N
DO 5 J=1,N
5 MI(I,J)=M(I,J)
CALL SIMUL (M1,ZA,N,0,LA,LB,S)
CALL EXTFUN (F,0.0,N)
DO 20 I=1,N
R(I)=F(I)

```

```

DO 10 J=1,N
10 R(I)=R(I)-C(I,J)*XD(I,J)-K(I,J)*X1(J)
20 CONTINUE
CALL XMULT (HI,R,XDDI,N)
DO 25 J=1,N
X(1,J)=X1(J)
XD(1,J)=XD(I,J)
25 XDD(1,J)=XDDI(J)
A1=6.0/((TH*DELTA)**2)
A2=3.0/(TH*DELTA)
A3=2.0*A2
A4=TH*DELTA/2.0
A5=A1/TH
A6=-A3/TH
A7=1.0-(3.0/TH)
A8=DELTA/2.0
A9=(DELTA**2)/6.0
DO 30 I=1,N
DO 30 J=1,N
30 TK(I,J)=K(I,J)+A1*M(I,J)+A2*C(I,J)
CALL SIMUL (TK,ZA,N,0,LA,LB,S)
DO 100 II=2,NSTEP1
DO 40 I=1,N
XN1(I)=A1*X(II-1,I)+A3*XD(II-1,I)+2.0*XDD(II-1,I)
40 XN2(I)=A2*X(II-1,I)+2.0*XD(II-1,I)+A4*XDD(II-1,I)
CALL XMULT (M,XN1,XN3,N)
CALL XMULT (C,XN2,XN4,N)
TIME1=REAL(II-2)*DELTA
TIME2=REAL(II-1)*DELTA
CALL EXTFUN (F1,TIME1,N)
CALL EXTFUN (F2,TIME2,N)
DO 45 J=1,N
45 FI(J)=F1(J)+TH*(F2(J)-F1(J))+XN3(J)+XN4(J)
CALL XMULT (TK,FT,XT,N)
DO 50 J=1,N
XDD(II,J)=A5*(XT(J)-X(II-1,J))+A6*XD(II-1,J)+A7*XDD(II-1,J)
XD(II,J)=XD(II-1,J)+A8*(XDD(II,J)+XDD(II-1,J))
50 X(II,J)=X(II-1,J)+DELTA*XD(II-1,J)+A9*(XDD(II,J)+2.0*XDD(II-1,J))
100 CONTINUE
RETURN
END

C =====
C
C SUBROUTINE EXTFUN
C
C =====
SUBROUTINE EXTFUN (F,TIME,N)
DIMENSION F(N)
F(1)=0.0
F(2)=10.0
RETURN
END

C =====
C
C SUBROUTINE XMULT
C
C =====
SUBROUTINE XMULT (A,B,BB,N)
DIMENSION A(N,N),B(N),BB(N)

```

```

      DO 10 I=1,N
      BB(I)=0.0
      DO 10 J=1,N
10    BB(I)=BB(I)+A(I,J)*B(J)
      RETURN
      END

```

SOLUTION BY WILSON METHOD

GIVEN DATA:

N= 2 NSTEP= 24 TH= 0.14000000E+01 DELTA= 0.24216267E+00

SOLUTION:

STEP	TIME	X(I,1)	XD(I,1)	XDD(I,1)	X(I,2)	XD(I,2)	XDD(I,2)
1	0.0000	0.3000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.5000E+01
2	0.2422	0.3344E-02	0.4143E-01	0.3422E+00	0.1392E+00	0.1119E+01	0.4244E+01
3	0.4843	0.2893E-01	0.1926E+00	0.9065E+00	0.5201E+00	0.1967E+01	0.2756E+01
4	0.7265	0.1072E+00	0.4742E+00	0.1419E+01	0.1058E+01	0.2395E+01	0.7809E+00
5	0.9687	0.2649E+00	0.8337E+00	0.1550E+01	0.1641E+01	0.2336E+01	-.1267E+01
6	1.2108	0.5076E+00	0.1152E+01	0.1076E+01	0.2153E+01	0.1825E+01	-.2958E+01
7	1.4530	0.8074E+00	0.1281E+01	-.4223E-02	0.2498E+01	0.9863E+00	-.3964E+01
8	1.6951	0.1104E+01	0.1106E+01	-.1442E+01	0.2619E+01	0.6096E-02	-.4131E+01
9	1.9373	0.1316E+01	0.5916E+00	-.2807E+01	0.2506E+01	-.9176E+00	-.3497E+01
10	2.1795	0.1369E+01	-.1866E+00	-.3621E+01	0.2193E+01	-.1615E+01	-.2266E+01
11	2.4216	0.1218E+01	-.1052E+01	-.3526E+01	0.1750E+01	-.1979E+01	-.7362E+00
12	2.6638	0.8710E+00	-.1772E+01	-.2420E+01	0.1264E+01	-.1974E+01	0.7743E+00
13	2.9060	0.3897E+00	-.2127E+01	-.5128E+00	0.8208E+00	-.1638E+01	0.2002E+01
14	3.1481	-.1186E+00	-.1981E+01	0.1717E+01	0.4905E+00	-.1059E+01	0.2781E+01
15	3.3903	-.5290E+00	-.1330E+01	0.3658E+01	0.3183E+00	-.3521E+00	0.3056E+01
16	3.6324	-.7333E+00	-.3125E+00	0.4747E+01	0.3207E+00	0.3643E+00	0.2860E+01
17	3.8746	-.6708E+00	0.8244E+00	0.4643E+01	0.4872E+00	0.9876E+00	0.2287E+01
18	4.1168	-.3477E+00	0.1791E+01	0.3340E+01	0.7852E+00	0.1440E+01	0.1451E+01
19	4.3589	0.1627E+00	0.2337E+01	0.1170E+01	0.1167E+01	0.1672E+01	0.4655E+00
20	4.6011	0.7389E+00	0.2323E+01	-.1288E+01	0.1575E+01	0.1660E+01	-.5639E+00
21	4.8433	0.1243E+01	0.1757E+01	-.3380E+01	0.1951E+01	0.1406E+01	-.1535E+01
22	5.0854	0.1558E+01	0.7952E+00	-.4566E+01	0.2239E+01	0.9371E+00	-.2339E+01
23	5.3276	0.1617E+01	-.3112E+00	-.4571E+01	0.2392E+01	0.3075E+00	-.2860E+01
24	5.5697	0.1418E+01	-.1283E+01	-.3457E+01	0.2382E+01	-.4004E+00	-.2987E+01
25	5.8119	0.1024E+01	-.1894E+01	-.1588E+01	0.2201E+01	-.1082E+01	-.2639E+01

11.50

```

C =====
C
C MAIN PROGRAM WHICH CALLS NUMARK
C
C =====
C FOLLOWING 10 LINES CONTAIN PROBLEM-DEPENDENT DATA
  REAL M(2,2),MI(2,2),K(2,2)
  DIMENSION C(2,2),XI(2),XDI(2),XDDI(2),X(25,2),XD(25,2),XDD(25,2),
  2 XT(2),F(2),R(2),LA(2),LB(2,2),S(2),ZA(2),
  3 TK(2,2),XN1(2),XN2(2),XN3(2),XN4(2)
  DATA N,NSTEP,NSTEP1,ALPHA,BETA,DELTA/2,24,25,0.16666667,0.5,
  2 0.24216267/
  DATA XI/0.0,0.0/
  DATA XDI/0.0,0.0/
  DATA M/1.0,0.0,0.0,2.0/
  DATA C/0.0,0.0,0.0,0.0/
  DATA K/6.0,-2.0,-2.0,8.0/
C END OF PROBLEM-DEPENDENT DATA

```

```

      CALL NUMARK (M,C,K,MI,XI,XDI,XDDI,X,XD,XDD,XT,F,R,LA,LB,S,ZA,
2 TK,XN1,XN2,XN3,XN4,NSTEP1,DELTA,ALPHA,BETA,N)
      PRINT 10
10  FORMAT (//,27H SOLUTION BY NEWMARK METHOD,/)
      PRINT 20, N,NSTEP,ALPHA,BETA,DELTA
20  FORMAT (12H GIVEN DATA:,,/ ,3H N=,15,4X,7H NSTEP=,15,4X,7H ALPHA=,
2 E15.8,2X,6H BETA=,E15.8,2X,7H DELTA=,E15.8,/)
      PRINT 30
30  FORMAT (10H SOLUTION:,,/ ,5H STEP,3X,5H TIME,3X,7H X(I,1),3X,
2 8H XD(I,1),2X,9H XDD(I,1),4X,7H X(I,2),3X,8H XD(I,2),2X,
3 9H XDD(I,2),/)
      DO 40 I=1,NSTEP1
      TIME=REAL(I-1)*DELTA
40  PRINT 50, I,TIME,X(I,1),XD(I,1),XDD(I,1),X(I,2),XD(I,2),XDD(I,2)
50  FORMAT (1X,I4,F8.4,6(1X,E10.4))
      STOP
      END

C =====
C
C SUBROUTINE NUMARK
C
C =====
      SUBROUTINE NUMARK (M,C,K,MI,XI,XDI,XDDI,X,XD,XDD,XT,F,R,LA,LB,S,
2 ZA,TK,XN1,XN2,XN3,XN4,NSTEP1,DELTA,ALPHA,BETA,N)
      REAL M(N,N),MI(N,N),K(N,N)
      DIMENSION C(N,N),XI(N),XDI(N),XDDI(N),X(NSTEP1,N),XD(NSTEP1,N),
2 XDD(NSTEP1,N),XT(N),F(N),R(N),LA(N),LB(N,2),
3 S(N),ZA(N),TK(N,N),XN1(N),XN2(N),XN3(N),XN4(N)
      DO 5 I=1,N
      DO 5 J=1,N
5  MI(I,J)=M(I,J)
      CALL SIMUL (MI,ZA,N,0,LA,LB,S)
      CALL EXTFUN (F,0.0,N)
      DO 20 I=1,N
      R(I)=F(I)
      DO 10 J=1,N
10  R(I)=R(I)-C(I,J)*XDI(J)-K(I,J)*XI(J)
20  CONTINUE
      CALL XMULT (MI,R,XDDI,N)
      DO 25 J=1,N
      X(1,J)=XI(J)
      XD(1,J)=XDI(J)
25  XDD(1,J)=XDDI(J)
      A1=1.0/(ALPHA*(DELTA**2))
      A2=1.0/(ALPHA*DELTA)
      A3=(1.0/(2.0*ALPHA))-1.0
      A4=(1.0-BETA)*DELTA
      A5=BETA*DELTA
      A6=BETA/(ALPHA*DELTA)
      A7=(BETA/ALPHA)-1.0
      A8=(A7-1.0)*DELTA/2.0
      DO 30 I=1,N
      DO 30 J=1,N
30  TK(I,J)=A1*M(I,J)+A6*C(I,J)+K(I,J)
      CALL SIMUL (TK,ZA,N,0,LA,LB,S)
      DO 100 II=2,NSTEP1

```

```

      TIME=REAL(II-1)*DELTA
      CALL EXTFUN (F,TIME,N)
      DO 40 J=1,N
      XN1(J)=A1*X(II-1,J)+A2*XD(II-1,J)+A3*XDD(II-1,J)
40    XN2(J)=A6*X(II-1,J)+A7*XD(II-1,J)+A8*XDD(II-1,J)
      CALL XMULT (M,XN1,XN3,N)
      CALL XMULT (C,XN2,XN4,N)
      DO 45 J=1,N
45    XN1(J)=F(J)+XN3(J)+XN4(J)
      CALL XMULT (TK,XN1,XT,N)
      DO 50 J=1,N
      X(II,J)=XT(J)
      XDD(II,J)=A1*(X(II,J)-X(II-1,J))-A2*XD(II-1,J)-A3*XDD(II-1,J)
50    XD(II,J)=XD(II-1,J)+A4*XDD(II-1,J)+A5*XDD(II,J)
100  CONTINUE
      RETURN
      END

```

```

C =====
C
C SUBROUTINE EXTFUN
C
C =====
      SUBROUTINE EXTFUN (F,TIME,N)
      DIMENSION F(N)
      F(1)=0.0
      F(2)=10.0
      RETURN
      END

```

```

C =====
C
C SUBROUTINE XMULT
C
C =====
      SUBROUTINE XMULT (A,B,BB,N)
      DIMENSION A(N,N),B(N),BB(N)
      DO 10 I=1,N
      BB(I)=0.0
      DO 10 J=1,N
10    BB(I)=BB(I)+A(I,J)*B(J)
      RETURN
      END

```

#### SOLUTION BY NEWMARK METHOD

##### GIVEN DATA:

N= 2      NSTEP= 24      ALPHA= 0.16666667E+00      BETA= 0.50000000E+00

##### SOLUTION:

DELTA= 0.24216267E+00

STEP	TIME	X(I,1)	XD(I,1)	XDD(I,1)	X(I,2)	XD(I,2)	XDD(I,2)
1	0.0000	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.5000E+01
2	0.2422	0.2606E-02	0.3228E-01	0.2666E+00	0.1411E+00	0.1143E+01	0.4438E+01
3	0.4843	0.2461E-01	0.1757E+00	0.9181E+00	0.5329E+00	0.2030E+01	0.2893E+01
4	0.7265	0.1005E+00	0.4775E+00	0.1574E+01	0.1088E+01	0.2471E+01	0.7467E+00
5	0.9687	0.2644E+00	0.8845E+00	0.1788E+01	0.1687E+01	0.2382E+01	-0.1483E+01

6	1.2108	0.5257E+00	0.1252E+01	0.1251E+01	0.2203E+01	0.1805E+01	-.3285E+01
7	1.4530	0.8530E+00	0.1398E+01	-.5063E-01	0.2534E+01	0.8884E+00	-.4281E+01
8	1.6951	0.1173E+01	0.1175E+01	-.1792E+01	0.2623E+01	-.1529E+00	-.4318E+01
9	1.9373	0.1389E+01	0.5460E+00	-.3400E+01	0.2467E+01	-.1097E+01	-.3481E+01
10	2.1795	0.1413E+01	-.3806E+00	-.4253E+01	0.2114E+01	-.1766E+01	-.2042E+01
11	2.4216	0.1200E+01	-.1369E+01	-.3914E+01	0.1643E+01	-.2058E+01	-.3706E+00
12	2.6638	0.7690E+00	-.2124E+01	-.2317E+01	0.1148E+01	-.1960E+01	0.1175E+01
13	2.9060	0.2111E+00	-.2384E+01	0.1724E+00	0.7195E+00	-.1536E+01	0.2333E+01
14	3.1481	-.3348E+00	-.2017E+01	0.2854E+01	0.4223E+00	-.8929E+00	0.2976E+01
15	3.3903	-.7196E+00	-.1078E+01	0.4907E+01	0.2946E+00	-.1570E+00	0.3102E+01
16	3.6324	-.8292E+00	0.2025E+00	0.5664E+01	0.3445E+00	0.5568E+00	0.2793E+01
17	3.8746	-.6221E+00	0.1475E+01	0.4843E+01	0.5550E+00	0.1156E+01	0.2158E+01
18	4.1168	-.1445E+00	0.2382E+01	0.2647E+01	0.8899E+00	0.1574E+01	0.1296E+01
19	4.3589	0.4811E+00	0.2667E+01	-.2884E+00	0.1299E+01	0.1766E+01	0.2842E+00
20	4.6011	0.1091E+01	0.2257E+01	-.3098E+01	0.1724E+01	0.1702E+01	-.8069E+00
21	4.8433	0.1529E+01	0.1281E+01	-.4966E+01	0.2103E+01	0.1377E+01	-.1862E+01
22	5.0854	0.1689E+01	0.2673E-01	-.5390E+01	0.2372E+01	0.8102E+00	-.2799E+01
23	5.3276	0.1548E+01	-.1150E+01	-.4326E+01	0.2480E+01	0.6281E-01	-.3374E+01
24	5.5697	0.1163E+01	-.1938E+01	-.2188E+01	0.2396E+01	-.7600E+00	-.3421E+01
25	5.8119	0.6543E+00	-.2166E+01	0.3099E+00	0.2118E+01	-.1515E+01	-.2817E+01

---