



Standard Practice for Sensory Evaluation of Edible Oils and Fats¹

This standard is issued under the fixed designation E1627; the number immediately following the designation indicates the year of original adoption or, in the case of revision, the year of last revision. A number in parentheses indicates the year of last reapproval. A superscript epsilon (ϵ) indicates an editorial change since the last revision or reapproval.

1. Scope

1.1 This practice covers the recommended procedures for the sensory evaluation of edible oils and fats.

1.2 This practice covers techniques for evaluating odor and flavor in fats and oils, for determining overall odor and flavor intensity, and the intensity of individual odors or flavors.

1.3 The techniques used in this practice are applicable to oils (liquid at room temperature) and liquified fats (solid at room temperature).

1.4 The values in SI units are to be regarded as the standard.

1.5 *This standard does not purport to address all of the safety concerns, if any, associated with its use. It is the responsibility of the user of this standard to establish appropriate safety and health practices and determine the applicability of regulatory limitations prior to use.* Specific precautions are given in Section 7.

2. Referenced Documents

2.1 *ASTM Standards:*²

E1346 Practice for Bulk Sampling, Handling, and Preparing Edible Vegetable Oils for Sensory Evaluation

3. Terminology

3.1 A lexicon specific for descriptors of odors and flavors in oils and fats is included in Appendix X2.

4. Summary of Practice

4.1 This practice addresses the procedures for screening and training of oil assessors; rating and scoring samples; and data collection, handling, analysis, and interpretation.

5. Significance and Use

5.1 The application of this practice will help ensure consistency in procedures used for the sensory evaluation of edible oils.

¹ This practice is under the jurisdiction of ASTM Committee E18 on Sensory Evaluation and is the direct responsibility of Subcommittee E18.06 on Food and Beverage Evaluation.

Current edition approved Aug. 1, 2011. Published August 2011. Originally approved in 1994. Last previous edition approved in 2004 as E1627 – 94 (2004). DOI: 10.1520/E1627-11.

² For referenced ASTM standards, visit the ASTM website, www.astm.org, or contact ASTM Customer Service at service@astm.org. For *Annual Book of ASTM Standards* volume information, refer to the standard's Document Summary page on the ASTM website.

5.2 This practice is designed for use by oil processors or research laboratories for evaluations by a trained, experienced sensory panel under the supervision of a sensory professional or for use by quality control and quality assurance personnel for the sensory evaluation of edible oils and fats.

6. Apparatus

6.1 *Glass Vial*, 30-mm outside diameter by 57-mm height, wide-mouth threaded top. Use amber glass for odor/flavor evaluations; clear glass for visual examination. Alternatively, use 2 oz sample cups also for odor/flavor evaluations only.

6.2 *Circulating Waterbath*, with automatic timer, thermostat, and rack.

6.3 *Waterbath Thermometer*, with range from 20 to 100°C in 1°C divisions, calibrated for 76-mm immersion, 305 mm long.

6.4 *Hard plastic threaded caps* with liners, or tape (PFTE pipe thread tape), to cover top of vial opening before capping with new, nonmetallic screw-type caps. Tape should completely cover vial opening or multiple strips of tape should be used.

7. Precautions

7.1 Assessors and sample servers should avoid introducing extraneous odors during testing by use of products, such as scented hand soap, hand creams, perfume, etc., or odorous writing instruments or inks. Assessors should avoid exhaling into sample vials.

8. Procedures for Recruitment and Screening Assessors

8.1 For basic information on conducting sensory tests, see MNL26³ and STP 758.⁴

8.2 For normal sensory acuity for basic tastes, see STP 758.⁴

8.3 *General Odor or Flavor Recognition Relating to Oils*—Present prospective assessors with a series of samples and a list of applicable oil descriptors specific for the type of oil to be tested (see Appendix X1). Appendix X2 contains definitions, reference standards, and examples of each descriptor. Test prospective assessors for general discrimination and the ability to describe samples and demonstrate familiarity with terms.

³ *Manual on Sensory Testing Methods*, ASTM MNL26, ASTM International.

⁴ *Guidelines for Selection and Training of Sensory Panel Members*, ASTM STP 758, ASTM International, 1981.

8.4 For general interest and availability, see STP 758.⁴

9. Procedures for Training Oil Assessors

9.1 See STP 758⁴ for information on panel training.

9.2 Determine training based upon test objective. Tests may include intensity ranking, attribute recognition, or difference tests, or a combination thereof (see MNL26³).

9.3 *Terminology/Characteristics (See 3.1):*

9.3.1 Examples of odor, flavors, and tastes predominately characteristic of various oil types are presented in **Appendix X2**. Attributes are identified as typical of an unprocessed or partially processed oil (U), freshly processed oil (F), deteriorated oil (D), or origin unknown (X). The appendix is a general guideline based on the attributes typically identified for each oil type; however, other attributes may be noted.

9.4 Prepare training samples characteristic of various odors or flavors and various intensity levels. Use **Appendix X1** and **Appendix X2** as guides.

9.5 Evaluate a series of concentrations starting with easily distinguished samples and proceed to more difficult discriminations.

9.6 Evaluate assessors' consistency on repeated tests as recommended in STP 758.⁴

10. Procedures for Oil Sample Handling, Preparation, and Presentation

10.1 For information on serving containers, sample size, heating methods, sample temperature, and presentation methods, see Practice **E1346**.

10.2 Oils should not be held at serving temperature for more than 60 min to prevent deterioration from oxidation.

10.3 If samples are presented in pairs or other multiples, it is recommended that a method be used to maintain uniform sample temperature of the oils during testing. Aluminum blocks, with recesses to hold vials, heated at a temperature of 5°C higher than the serving temperature of the oil will keep the sample at the proper serving temperature for 10 min. Molded styrene (styrofoam) blocks, with recesses to hold vials, will help minimize temperature loss. Vials should fit into the recesses or cavities in the blocks deep enough so the oil line in the vial does not extend above the top of the recess. The diameter of the aluminum block recess should not be more than 1 cm wider than the diameter of the vial to allow adequate heat transfer.

11. Instructions to Assessors for Odor Evaluations

11.1 Evaluate the oils for odor in the order presented from left to right.

11.2 Pick up the vial containing the oil; hold the vial as close to base as possible.

11.3 Swirl the covered vial; lift to nose; remove the cover; sniff the headspace above the oil (use short, "bunny" sniffs); replace the cover quickly.

11.4 Sniff in the same manner—distance from nose, number of times, length of time—for each sample.

11.5 Smell back of hand before testing samples and between samples to help "zero" your nose and to prevent adaptation to oil odors.

11.6 If testing oils with weak odors, smell an empty container to facilitate adaptation to extraneous odors and to allow for better discrimination between oils.

12. Instructions to Assessors for Flavor Evaluations

12.1 Rinse mouth well with warm filtered water ($50 \pm 1^\circ\text{C}$) before starting the flavor evaluation.

12.2 Taste the samples in the order presented from left to right.

12.3 Put the entire 10-mL sample of warm oil into the mouth; swish through the mouth thoroughly; cup mouth and draw air in through mouth and exhale through nose to enhance perception of aromatics.

12.4 Expectorate the sample; do not swallow the oil.

12.5 Rinse the mouth well with warm water ($50 \pm 1^\circ\text{C}$) between samples for a predetermined amount of time to clear mouth of residual flavors.

12.6 Wait a predetermined amount of time before tasting subsequent samples to prevent taste fatigue; be consistent.

12.7 Additional methods to clear the mouth include unsalted soda crackers, 50:50 blend of warm water and sodium-free carbonated water ($50 \pm 1^\circ\text{C}$).

12.8 If residual flavors persist, repeat the procedure of rinsing and resting.

13. Procedures for Data Collection

13.1 Discrimination tests, for example, Triangle, Duo-trio, A not A, etc., are used to determine if a difference exists between two samples. Uses include qualifying alternate ingredient suppliers; confirming quality control in the plant, determining end of shelf-life; and reformulation of existing brands (see Chapter 2 of MNL26³).

13.2 Descriptive or scalar scoring tests are used to rate the overall intensity of a sample and to describe characteristic odors and flavors of samples. Use to find sensory differences between competitive products, aged products, new formulas, etc., and to interpret results of consumer tests and understand the effects of technical variables on product attributes (see MNL 13⁵).

13.3 Quality tests are used to rate the overall quality of a fat or oil with moderate to strong characteristic flavors such as olive or peanut oil (see MNL26³).

14. Data Handling

14.1 Statistical analysis of the data will depend on the type of test and test design. MNL26³ contains statistical analysis appropriate for various sensory tests. Data handling methods for descriptive tests are presented in MNL 13.⁵

15. Data Interpretation

15.1 Action criteria will depend on the policy of the laboratory or company and will be product specific. Policies will determine the intensity levels of specific flavors that are desired or will be permitted. The intensities allowed will vary based on the attribute and its positive or negative contribution to the oil or fat. Customer complaints will validate decisions over time.

⁵ *Manual on Descriptive Analysis Testing, ASTM MNL 13*, ASTM, 1992.

16. Keywords

16.1 descriptive testing; discriminative testing; flavor; odor; scaling; sensory analysis; taste

APPENDIXES

(Nonmandatory Information)

X1. VOCABULARY AND OIL ATTRIBUTES CHARACTERISTIC OF UNPROCESSED OIL (U), FRESHLY PROCESSED OIL (F), DETERIORATED OIL (D), OR ORIGIN UNIDENTIFIED (X)

TABLE X1.1 Oil Types and Attributes^A

Attributes	Oil Type												
	Corn	Cotton Seed	Coconut	Fish/ Marine	MCT ^B	Olive	Palm	Peanut	Canola Rapeseed	Ricebran	Safflower	Soy	Sunflower
Bacony	...	...	U	...	...	...	X	...	...	...	...	...	...
Beany	...	...	...	...	...	...	...	...	D	...	...	D/U	...
Bitter	...	...	...	...	...	...	X	...	...	...	...	...	...
Burnt	U	...	X	...	...	...	...	X	...	X	...	U	X
Buttery	F	F	F	...	F	...	F	X	F	...	F	F	F
Cardboard	D	...	...	...	...	...	D	D	...	...	...	D	D
Corny	F	...	...	...	...	...	...	...	...	...	...	...	...
Fishy	...	...	...	U/F/D	...	...	...	...	D	...	...	D	...
Fruity	...	...	...	...	...	F	...	F	...	...	...	...	...
Grassy	X	...	...	...	...	...	...	X	D	X	X	U/D	U/D
Green	...	...	...	...	...	X	...	...	D	X	...	...	...
Hay	X	...	...	...	...	...	...	X	...	X	X	U/D	U/D
Hully	...	...	...	...	...	...	...	...	...	...	...	...	X
Nutty	F	X	X	...	X	...	X	F	...	X	...	F	...
Painty	...	...	...	...	...	...	...	...	D	...	...	D	...
Pine	...	...	...	...	...	...	...	...	...	...	...	...	U
Rancid	D	D	D	...	...	D	D	D	D	D	D	D	D
Rubbery	X	...	...	...	...	...	...	...	X	...	...	X	...
Soapy	...	...	X	...	...	...	X	...	...	...	...	...	...
Sulfur	...	...	...	...	...	...	...	...	U/D	...	...	...	...
Waxy	X	...	X	...	U	...	...	...	...	X	X	...	X
Weedy	X	...	...	...	...	...	...	X	X	X	X	X	X
Woody	...	...	...	...	...	...	...	X	...	X	X	...	X

^A U = characteristic of unprocessed or partially processed oil

F = characteristic of freshly processed oil

D = characteristic of deteriorated oil.

X = unidentified origin.

Other flavors may be present from contamination, processing conditions, etc.: pumpkin, melon, watermelon, petroleum, metallic, musty.

^B Medium chain triglycerides.

X2. LEXICON FOR FATS AND OILS

Bacon		<i>Definition—</i>	An aromatic reminiscent of smoked bacon.	<i>Definition—</i>	An aromatic reminiscent of burnt popcorn or grains.
<i>Reference—</i>	Crude undeodorized coconut oil heated to 38°C.	<i>Reference—</i>		<i>Reference—</i>	Crude, unprocessed corn oil.
<i>Example—</i>	Fried smoked pork bacon.	<i>Example—</i>		<i>Example—</i>	Air-popped popcorn.
Beany		<i>Definition—</i>	An aromatic characteristic of raw soybeans.	<i>Definition—</i>	An aromatic reminiscent of fresh, sweet, unsalted butter.
<i>Reference—</i>	Crude soybean oil diluted in fresh soybean oil (5:95).	<i>Reference—</i>		<i>Reference—</i>	Fresh, sweet, unsalted butter diluted in good quality soybean oil (1:99).
<i>Example—</i>	Ground lima beans (dry mixed with water (2:98 ratio)).	<i>Example—</i>		<i>Example—</i>	Freshly processed unsalted butter.
Bitter		<i>Definition—</i>	A basic taste simulated by such substances as quinine and caffeine.	<i>Definition—</i>	An aromatic associated with the odor of wet cardboard or paper.
<i>Reference—</i>	0.2 % caffeine in water.	<i>Reference—</i>		<i>Reference—</i>	Wet one cup unsalted, dry-roasted vacuum-packed peanuts with distilled water; place wet nuts on tray to air-dry for 24 h.
<i>Example—</i>	Tonic water.	<i>Example—</i>		<i>Example—</i>	Wet cardboard.
Bland		<i>Definition—</i>	No aromatics or taste factors perceptible.	<i>Definition—</i>	
<i>Example—</i>	Mineral oil.	<i>Example—</i>		<i>Example—</i>	
Burnt		<i>Definition—</i>		<i>Definition—</i>	An aromatic of steeped ground corn.



<i>Reference—</i>	Crude corn oil diluted in fresh corn oil (5:95).	<i>Painty</i>	
<i>Example—</i>	Raw corn: non-heat-treated corn; cooked corn: heated or boiled corn; and, toasted corn: heated enough to caramelize sugars.	<i>Definition—</i>	An aromatic reminiscent of oils containing linolenic acid such as linseed or rapeseed (canola) oil; not noted in non-linolenic acid oils such as peanut.
<i>Fishy</i>		<i>Reference—</i>	Good-quality canola oil aged for four to eight days at 60°C or until a peroxide value of 10.0 is reached.
<i>Definition—</i>	An aromatic reminiscent of cod liver oil.	<i>Example—</i>	Linseed oil.
<i>Reference—</i>	Cod liver oil diluted in good-quality soybean oil (1:99).	<i>Pine</i>	
<i>Example—</i>	Odor from canola (rapeseed) oil heated at 190°C.	<i>Definition—</i>	An aromatic reminiscent of pine needles noted in sunflower oil.
<i>Fruity</i>		<i>Reference—</i>	Bleached, undeodorized sunflower oil diluted (5:95) in good-quality fresh soybean oil.
<i>Definition—</i>	An aromatic reminiscent of ripe fruit.	<i>Example—</i>	Fresh pine needles cut in small pieces.
<i>Reference—</i>	2 ppm ethyl acetate.	<i>Plastic</i>	
<i>Example—</i>	Olive oil.	<i>Definition—</i>	An aromatic reminiscent of plastic containers or food stored in a plastic container.
<i>Grassy</i>		<i>Reference—</i>	Plastic strips from a poly(ethylene terephthalate) (PET) package stored 24 h in fresh, good-quality soybean oil.
<i>Definition—</i>	An aromatic reminiscent of the green character of mowed grass.	<i>Example—</i>	Plastic cup.
<i>Reference—</i>	Crude soybean oil from non-heat-treated soybeans diluted in good-quality soybean oil (5:95).	<i>Rancid</i>	
<i>Example—</i>	Fresh cut grass.	<i>Definition—</i>	An aromatic reminiscent of odor or flavor of highly oxidized oils containing high amounts of linoleic acid such as sunflower, cottonseed, or peanut.
<i>Green</i>		<i>Reference—</i>	Good-quality cottonseed oil aged for four days at 60°C or until a peroxide value of approximately 5.0 is reached.
<i>Definition—</i>	An aromatic associated with unprocessed immature fruits or grains.	<i>Example—</i>	Potato chips fried in cottonseed oil and aged.
<i>Reference—</i>	5 ppm cis-3-hexenol in water.	<i>Reverted</i>	
<i>Example—</i>	Raw immature soybeans.	<i>Definition—</i>	A general term denoting the process of oil flavor reversion. During initial stages of oxidation, most oils develop flavors or odors reminiscent of odors or flavors detectable in crude oil. Each oil type would have a characteristic reverted odor or flavor. Not recommended as a specific odor or flavor.
<i>Hay</i>		<i>Rubbery</i>	
<i>Definition—</i>	An aromatic reminiscent of dried grass character of air-dried grain or vegetation.	<i>Definition—</i>	An aromatic reminiscent of old rubber.
<i>Reference—</i>	Crude soybean oil from heat-treated beans diluted in good-quality soybean oil (5:95).	<i>Reference—</i>	0.5 ppm methyl allyl trisulfide (odor only).
<i>Example—</i>	Dried alfalfa.	<i>Example—</i>	Poorly processed corn oil; rubber stoppers.
<i>Hully</i>		<i>Soapy</i>	
<i>Definition—</i>	An aromatic associated with the outer protective coating of a grain or oilseed.	<i>Definition—</i>	An aromatic reminiscent of unscented soap.
<i>Reference—</i>	Raw steeped peanut hulls.	<i>Reference—</i>	Ivory brand unscented soap flakes.
<i>Example—</i>	Sunflower hulls (confectionery type).	<i>Example—</i>	Oxidized fat containing lauric acid, such as coconut oil.
<i>Hydrogenated</i>		<i>Sulfur</i>	
<i>Definition—</i>	An aromatic reminiscent of the sweet paraffin-like odor of crayons.	<i>Definition—</i>	An aromatic reminiscent of oils from seeds in the sulfur-containing vegetable family such as rapeseed (canola).
<i>Reference—</i>	10 % undeodorized hydrogenated soybean oil (iodine value = 90–110) in good-quality soybean oil.	<i>Reference—</i>	Bleached, undeodorized canola oil diluted in good-quality canola oil (5:95).
<i>Example—</i>	All vegetable solid shortening.	<i>Example—</i>	Brussels sprouts.
<i>Light-struck</i>		<i>Waxy</i>	
<i>Definition—</i>	Mixture of aromatics characteristic of light-sensitive oils such as soybean that are exposed to fluorescent light or sunlight.	<i>Definition—</i>	An aromatic reminiscent of candle wax.
<i>Reference—</i>	Good-quality soybean oil exposed to fluorescent light (100 footcandles for one week or 800 footcandles for 4 h).	<i>Reference—</i>	High oleic sunflower oil heated to 190°C for 30 min.
<i>Melon</i>		<i>Example—</i>	Melted paraffin.
<i>Definition—</i>	An aromatic reminiscent of watermelon rind.	<i>Weedy</i>	
<i>Reference—</i>	0.002 ppm 2,6-nonadienal in good-quality soybean oil (odor only).	<i>Definition—</i>	An aromatic reminiscent of freshly cut weeds.
<i>Example—</i>	Soybean oil processed with phosphoric acid; watermelon rind.	<i>Reference—</i>	10 000 ppm 2-isobutylthiazole in propylene glycol.
<i>Metallic</i>		<i>Example—</i>	Mixture of freshly cut green weeds.
<i>Definition—</i>	An aromatic associated with metal coins.	<i>Woody</i>	
<i>Reference—</i>	0.01 % ferrous sulfate diluted in distilled, filtered water.	<i>Definition—</i>	An aromatic reminiscent of fresh, dry cut wood.
<i>Example—</i>	Copper pennies soaked in filtered water for 12 h; soybean oil processed without citric acid.	<i>Reference—</i>	Wood (oak) chips.
<i>Musty</i>		<i>Example—</i>	Peanut oil.
<i>Definition—</i>	An aromatic reminiscent of odor of a moldy or damp cellar or room.		
<i>Reference—</i>	25 ppb methyl isoborenol.		
<i>Example—</i>	Damp cloth stored in a plastic bag.		
<i>Nutty</i>			
<i>Definition—</i>	An aromatic reminiscent of fresh, sweet nutmeats.		
<i>Reference—</i>	Freshly ground English walnuts.		
<i>Example—</i>	Freshly processed peanut oil.		
<i>Oxidized</i>			
<i>Definition—</i>	A general term denoting the process of oxidative deterioration of oil. Oxidized flavors or odors range widely from buttery, grassy, rancid, to painty. Not recommended as a specific odor or flavor.		

ASTM International takes no position respecting the validity of any patent rights asserted in connection with any item mentioned in this standard. Users of this standard are expressly advised that determination of the validity of any such patent rights, and the risk of infringement of such rights, are entirely their own responsibility.

This standard is subject to revision at any time by the responsible technical committee and must be reviewed every five years and if not revised, either reapproved or withdrawn. Your comments are invited either for revision of this standard or for additional standards and should be addressed to ASTM International Headquarters. Your comments will receive careful consideration at a meeting of the responsible technical committee, which you may attend. If you feel that your comments have not received a fair hearing you should make your views known to the ASTM Committee on Standards, at the address shown below.

This standard is copyrighted by ASTM International, 100 Barr Harbor Drive, PO Box C700, West Conshohocken, PA 19428-2959, United States. Individual reprints (single or multiple copies) of this standard may be obtained by contacting ASTM at the above address or at 610-832-9585 (phone), 610-832-9555 (fax), or service@astm.org (e-mail); or through the ASTM website (www.astm.org). Permission rights to photocopy the standard may also be secured from the ASTM website (www.astm.org/COPYRIGHT/).