



Standard Guide for Sampling Seized Drugs for Qualitative and Quantitative Analysis¹

This standard is issued under the fixed designation E 2548; the number immediately following the designation indicates the year of original adoption or, in the case of revision, the year of last revision. A number in parentheses indicates the year of last reapproval. A superscript epsilon (ε) indicates an editorial change since the last revision or reapproval.

1. Scope

1.1 This guide covers minimum considerations for sampling of seized drugs for qualitative and quantitative analysis.

2. Referenced Documents

2.1 ASTM Standards:²

E 105 Practice for Probability Sampling Of Materials

E 122 Practice for Calculating Sample Size to Estimate, With a Specified Tolerable Error, the Average for a Characteristic of a Lot or Process

E 141 Practice for Acceptance of Evidence Based on the Results of Probability Sampling

E 1732 Terminology Relating to Forensic Science

E 2329 Practice for Identification of Seized Drugs

E 2334 Practice for Setting an Upper Confidence Bound For a Fraction or Number of Non-Conforming items, or a Rate of Occurrence for Non-conformities, Using Attribute Data, When There is a Zero Response in the Sample

2.2 ISO Standards:³

ISO 3534-1 Statistics – Vocabulary and symbols – Part 1: Probability and general statistical terms

ISO 3534-2 Statistics – Vocabulary and symbols – Part 2: Statistical quality control

3. Significance and Use

3.1 This guide provides information for the sampling of seized-drug submissions.

3.2 The principal purpose of sampling in the context of this guide is to answer relevant questions about a population by examination of a portion of the population. For example:

What is the net weight of the population?

What portion of the units of a population can be said to contain a given drug at a given level of confidence?

3.3 By developing a sampling strategy and implementing appropriate sampling schemes, as illustrated in Fig. 1, a laboratory will minimize the total number of required analytical determinations, while ensuring that all relevant legal and scientific requirements are met.

4. Sampling Strategy

4.1 A sampling strategy is highly dependent on the purpose of the investigation, the original question, and the ultimate use of the results. Laws and legal practices form the foundation of most strategies and shall be taken into account when designing a sampling scheme. Therefore, specific sampling strategies are not defined in this guide.

4.2 The laboratory has the responsibility to develop its own strategies consistent with these recommendations. It is recommended that the following key points be addressed:

4.2.1 Sampling may be statistical or non-statistical.

NOTE 1—For the purpose of this guide, the use of the term *statistical* is meant to include the notion of an approach that is probability-based.

4.2.1.1 In many cases, a non-statistical approach may suffice. The sampling plan shall provide an adequate basis for answering questions of applicable law. For example,

Is there a drug present in the population?

Are statutory enhancement levels satisfied by the analysis of a specified number of units?

4.2.1.2 If an inference about the whole population is to be drawn from a sample, then the plan shall be statistically based and limits of the inference shall be documented.

4.2.2 Statistically selected units shall be analyzed to meet Practice **E 2329** if statistical inferences are to be made about the whole population.

5. Sampling Scheme

5.1 The sampling scheme is an overall approach that includes population determination, selection of the sampling plan and procedure and, when appropriate, sample reduction prior to analysis (Fig. 2).

5.2 *Population Determination:*

¹ This guide is under the jurisdiction of ASTM Committee E30 on Forensic Sciences and is the direct responsibility of Subcommittee E30.01 on Criminalistics. Current edition approved March 1, 2007. Published April 2007.

² For referenced ASTM standards, visit the ASTM website, www.astm.org, or contact ASTM Customer Service at service@astm.org. For *Annual Book of ASTM Standards* volume information, refer to the standard's Document Summary page on the ASTM website.

³ Available from International Organization for Standardization (ISO), 1 rue de Varembe, Case postale 56, CH-1211, Geneva 20, Switzerland, <http://www.iso.ch>.

Sampling Strategy — Based upon jurisdictional requirements




FIG. 1 Relationship of Various Levels Required in Sampling




FIG. 2 Example of a Sampling Scheme—A Decision Flowchart

5.2.1 The population determination shall take into account all typical forms and quantities in which exhibits may appear.

5.2.2 A population can consist of a single unit or multiple units.

5.2.3 A multiple unit population shall consist of items that are similar in relevant visual characteristics.

Sampling Plan

5.3 There are numerous sampling plans used in the forensic analysis of drugs that are applicable to single and multiple unit populations.

5.4 When a single unit or bulk population is to be analyzed, the issue of homogeneity shall be addressed within the sampling plan.

5.4.1 One sample is sufficient if the bulk material is homogeneous. Analysts can make bulk material homogeneous.

5.4.2 If the bulk material is not homogeneous, several samples from different locations may be necessary to ensure that the test results are representative of the bulk material and to avoid false negative results.

5.5 Depending upon the inference to be drawn from the analysis for a multiple unit population, the sampling plan may be statistical or non-statistical.

5.5.1 Statistical approaches are applicable when inferences are made about the whole population. For example:

The probability that a given percentage of the population contains the drug of interest or is positive for a given characteristic.

The total net weight of the population is to be extrapolated from the weight of a sample.

5.5.1.1 Published examples of statistical approaches involving general considerations:

- (1) Practice E 105.
- (2) Practice E 141.
- (3) Terminology E 1732.
- (4) *Guidelines on Representative Drug Sampling*.⁴
- (5) ISO 3534-1.
- (6) ISO 3534-2.

5.5.1.2 Published examples of statistical approaches involving the hypergeometric, normal, and other distributions from a frequentist perspective:

- (1) Frank, et al, *Journal of Forensic Sciences*.⁵
- (2) *Guidelines on Representative Drug Sampling*.⁴
- (3) Practice E 2334.
- (4) Practice E 122.

5.5.1.3 Published examples of statistical approaches involving the hypergeometric, normal, and other distributions from a Bayesian perspective:

- (1) Coulson, et al, *Journal of Forensic Sciences*.⁶
- (2) *Guidelines on Representative Drug Sampling*.⁴

5.5.2 Non-statistical approaches are appropriate if no inference is to be made about the whole population.

5.5.2.1 Published examples:

(1) The “square root method.”⁷

(2) Methods listed in “Arbitrary Sampling” in *Guidelines on Representative Drug Sampling*.⁴

5.5.2.2 Selection of a single unit from a multiple unit population may be appropriate under certain circumstances (for example, management directives and legislative or judicial requirements, or both).

5.6 *Sampling Procedure:*

5.6.1 Establish the procedure for selecting the number of units that will comprise a sample.

5.6.1.1 For non-statistical approaches, select a sample appropriate for the analytical objectives.

5.6.1.2 For statistical approaches it is recommended that random sampling be conducted.

5.6.2 *Select a Random Sample:*

5.6.2.1 A random sample is one selected without bias. Computer generated random numbers or random number tables are commonly employed for such tasks and these should be included in the sampling plan.

5.6.2.2 Random sampling of items using random number tables may not be practical in all cases. In these instances, an alternate sampling plan shall be designed and documented to approach random selection. A practical solution involves a “black box” method, which refers to one that will prevent the sampler from consciously selecting a specific item from the population (that is, all units are placed in a box and the samples for testing are selected without bias). Random sampling is discussed in the following references:

- (1) Practice E 105.
- (2) Frank, et al, *Journal of Forensic Sciences*.⁵
- (3) ISO 3534-1.
- (4) ISO 3534-2.

5.7 *Sample Reduction*—Sample reduction may be applied in cases where the weight or volume of the selected units is too large for laboratory analysis (Fig. 2, insert A).

6. Analysis

6.1 *Statistically Selected Sample(s)*—In accordance with 4.2.2, it is recommended that each unit comprising the sample be analyzed to meet Practice E 2329 if statistical inferences are to be made about the whole population.

6.2 *Non-statistically Selected Sample(s)*—Practice E 2329 shall be applied to at least one unit of the sample.

7. Documentation

7.1 Inferences based on use of a sampling plan and concomitant analysis shall be documented.

8. Keywords

8.1 analytical method; qualitative method; quantitative method; sampling; seized drug analytical method

⁴ *Guidelines on Representative Drug Sampling*, European Network of Forensic Science Institutes (ENFSI), 2004, (www.ENFSI.org).

⁵ “Representative Sampling of Drug Seizures in Multiple Containers,” Frank et al, *Journal of Forensic Sciences*, 1991, 36(2), pp. 350-357.

⁶ “How Many Samples from a Drug Seizure Need to Be Analyzed?” Coulson et al, *Journal of Forensic Sciences*, 2001, 46(6), pp. 1456-1461.

⁷ *Recommended Methods for Testing Opium, Morphine and Heroin: Manual for Use by National Drug Testing Laboratories*, United Nations Office on Drugs and Crime, 1998.

ASTM International takes no position respecting the validity of any patent rights asserted in connection with any item mentioned in this standard. Users of this standard are expressly advised that determination of the validity of any such patent rights, and the risk of infringement of such rights, are entirely their own responsibility.

This standard is subject to revision at any time by the responsible technical committee and must be reviewed every five years and if not revised, either reapproved or withdrawn. Your comments are invited either for revision of this standard or for additional standards and should be addressed to ASTM International Headquarters. Your comments will receive careful consideration at a meeting of the responsible technical committee, which you may attend. If you feel that your comments have not received a fair hearing you should make your views known to the ASTM Committee on Standards, at the address shown below.

This standard is copyrighted by ASTM International, 100 Barr Harbor Drive, PO Box C700, West Conshohocken, PA 19428-2959, United States. Individual reprints (single or multiple copies) of this standard may be obtained by contacting ASTM at the above address or at 610-832-9585 (phone), 610-832-9555 (fax), or service@astm.org (e-mail); or through the ASTM website (www.astm.org).